

Table Of Content

Journal Cover	2
Author[s] Statement	3
Editorial Team	4
Article information	5
Check this article update (crossmark)	5
Check this article impact	5
Cite this article	5
Title page	6
Article Title	6
Author information	6
Abstract	6
Article content	7

Academia Open



By Universitas Muhammadiyah Sidoarjo

Originality Statement

The author[s] declare that this article is their own work and to the best of their knowledge it contains no materials previously published or written by another person, or substantial proportions of material which have been accepted for the published of any other published materials, except where due acknowledgement is made in the article. Any contribution made to the research by others, with whom author[s] have work, is explicitly acknowledged in the article.

Conflict of Interest Statement

The author[s] declare that this article was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright Statement

Copyright © Author(s). This article is published under the Creative Commons Attribution (CC BY 4.0) licence. Anyone may reproduce, distribute, translate and create derivative works of this article (for both commercial and non-commercial purposes), subject to full attribution to the original publication and authors. The full terms of this licence may be seen at <http://creativecommons.org/licences/by/4.0/legalcode>

EDITORIAL TEAM

Editor in Chief

Mochammad Tanzil Multazam, Universitas Muhammadiyah Sidoarjo, Indonesia

Managing Editor

Bobur Sobirov, Samarkand Institute of Economics and Service, Uzbekistan

Editors

Fika Megawati, Universitas Muhammadiyah Sidoarjo, Indonesia

Mahardika Darmawan Kusuma Wardana, Universitas Muhammadiyah Sidoarjo, Indonesia

Wiwit Wahyu Wijayanti, Universitas Muhammadiyah Sidoarjo, Indonesia

Farkhod Abdurakhmonov, Silk Road International Tourism University, Uzbekistan

Dr. Hindarto, Universitas Muhammadiyah Sidoarjo, Indonesia

Evi Rinata, Universitas Muhammadiyah Sidoarjo, Indonesia

M Faisal Amir, Universitas Muhammadiyah Sidoarjo, Indonesia

Dr. Hana Catur Wahyuni, Universitas Muhammadiyah Sidoarjo, Indonesia

Complete list of editorial team ([link](#))

Complete list of indexing services for this journal ([link](#))

How to submit to this journal ([link](#))

Article information

Check this article update (crossmark)



Check this article impact (*)



Save this article to Mendeley



(*) Time for indexing process is various, depends on indexing database platform

Assessment Awareness about Hand Hygiene among Health Care Providers in Hospitals at Al_Nasiriyah City

Penilaian Kesadaran tentang Kebersihan Tangan di antara Penyedia Layanan Kesehatan di Rumah Sakit di Kota Al_Nasiriyah

Teeba T. Khudair, teebathamer@utq.edu.iq, (1)

Department of Pharmaceutical Science, College of Pharmacy, University of Thi-Qar, Thi-Qar, 64001, Iraq, Iraq

⁽¹⁾ Corresponding author

Abstract

Background: The World Health Organization emphasizes hand hygiene as a crucial method for controlling hospital infections, emphasizing the need for healthcare workers to have current knowledge on this topic.. Objective: The study evaluates hand hygiene awareness among healthcare providers in AL_Nasiriyah hospitals, identifying factors influencing HH abidance, crucial for patient safety and reducing infectious disease transmission. Methods: A cross-sectional study was conducted in various hospitals in the governorate of AL_Nasiriyah city during the period from May to July 2024 to evaluate awareness Associated to (HH) hygiene among health care providers.by using a special format which included two parts, the first part related to their demographic characteristic and the second part contain multiple questions A purposive sample is selected for the study which sample consisting (171) working various specializations .Date were gathered through the interviewed in workplace and analyzed by using spss 20 through the application of descriptive statistical included (frequencies, percentages, M.S,and S.D). The overall result: The study revealed that over thirty-two percent of the population (63.2%) and those with over six years of health experience have the highest awareness of hand hygiene, influenced by education and experience. Conclusion: This study shows a good level of awareness of hand hygiene among health care practitioners at mean score (1.39) level of awareness regarding answering the questionnaire information. Recommendations: The study suggests continuous training for competent authorities, provision of basic hand hygiene environments, alcohol and sterilizers, and recent hygiene items for health places.

Published date: 2024-10-31 00:00:00

Introduction

A fundamental aspect of effective infection control in hospital is adequate, suitable hand-hygiene practices in all Medical practitioners who are in direct and indirect contact with patients or patient's environment. In 2009, the National Hand Hygiene Initiative (NHHI) was Conducted in In hospitals nationwide) Australia (Objective was to enhancing hand_ hygiene practices and diminished health care associated Contagions [1-3].

In the hospital where this assessment was performed, a significant decrease in infection rates from *Staphylococcus aureus* Was attained. From (2013--2017), there was a decline in cases. Results however show the Minimum rates of hand hygiene practices Engagement persists among non-clinical personnel [4].

It's imperative to Priority on hand hygiene, within all staff members in hospitals, As a result of a major cause of Health complications and mortality in Hospital-acquired infection [5]. In 2015, there were Several cases of resistant bacteria have emerged in healthcare settings. The most effective preventive measure for infection prevention in various health care settings, including nursing homes is (antiseptic) hand rubbing [6-7]. This term Indicates to the application of Aseptic agent)hand rub(to Mitigate or suppress the growth of microorganisms Absent the need for an external source of water and requiring no washing or drying with towels or other devices. [8]. Previous research has signified to individual knowledge deficiencies Impact on Risk-free hand hygiene practices such as correct periods of hand washing and shortfalls in hand rub advisories. [9-10]. Individual influences as knowledge of the five moments of hand hygiene, behaviors including not wearing hand Accessories while nursing, and Implementing their learnings training, to improve requirement of (hand hygiene), are essential and basic criteria for infection safeguarding [11] .The hygiene courses and training could capably raise the comply to hand rubbing and minimize Infections in healthcare settings. Also effective HH demands adequate organizational factors including providing of hand rub, protective clothing, and strong indigenous efforts from the nursing management such as role modelling, According to Merton's concept of a role model, individuals who set a positive example and are worthy of imitation can influence others. Previous research has indicated that hand hygiene was performed more frequently when team members in higher hierarchical positions sanitized, serving as role models for others [12-13]. While a large proportion of multidrug-resistant infections in nursing homes could be avoided through appropriate hand hygiene behavior of nurses, this behavior is influenced by organizational factors such as hygiene training, availability of resources and improved role modelling of nursing managers [14].

Methods

Design of the Study

A descriptive design study in which assessment approach is applied to achieve the objectives of the study and was conducted Assessment of Health Care Providers Knowledge about hand hygiene in health center from the various hospitals of AL_Nasiriyah. questionnaire has been adopted to meet and achieve the objectives of the study. The questionnaire consists of two parts; part one demographic characteristics and part two consist of medical staff knowledge toward hand Hygiene behavior and its necessity practice scales. Validity is concerned with the extent to which an instrument corresponds. The content validity of instrument was established through a panel of (3) experts from different specialties. The results of the review of the questionnaire by the experts revealed that all of the experts agree that (17) items of the study instrument are clear and adequate for the measurement of the phenomenon. In order to determine the reliability of the study instrument. The pilot study was conducted in Al-Nasiriyah center in Thi-Qar governorate. It was carried out on five of healthcare workers. The study was executed from the period of May to July 2024. The Purpose of pilot study to confirm the clarity and content adequacy of the instrument structure throughout the subjects understanding and to determine the required modifications. To enhance the reliability of the questionnaire. To test the feasibility and to determine the average time required for answering the questionnaire.

Reliability of the Questionnaire:

1. The internal consistency of the instrument was determined through the
2. computation of Alpha Correlation Coefficient (Cronbach's Alpha). The Coefficient
3. Alpha was applied to determine the reliability of the present study instrument by
4. application of Statistical Package for Social Science Program (SPSS) version 20.0.

The Sample of the Study:

A purposive, non-probability sample of Health Care Providers (171) who work at medical department were selected based on the study criteria, and after obtaining a consent from them.

Inclusion Criteria for Selecting the Sample

The sample was selected according to the following criteria:

1. Those who work in medical departments
2. Both male and female genders should participate in the study.

Limitations of the Study

The generalizability of our findings is not known despite the diversity of health care workers' units and we did not assess hygiene technically and did not include night shift care workers who might have responded differently and we were unable to control for the possibility of a burden variable the work

Result and Discussion

Result

		Frequency	Percent
Age	20-25	28	16.4
	26-30	35	20.5
	>30	108	63.2
	Total	171	100.0
Gander	Male	128	74.9
	Female	43	25.1
	Total	171	100.0
Level of qualification	Preparatory	38	22.2
	Institute	70	40.9
	College	51	29.8
	High studies	12	7.1
	Total	171	100.0
Years of experiences in hospital	1-2	21	12.3
	3-5	25	14.6
	>6	125	73.1
	Total	171	100.0
Marital status	Single	41	24.0
	Married	126	73.7
	Others	4	2.3
	Total	171	100.0
Residence	City	152	88.9
	Village	19	11.1
	Total	171	100.0

Table 1. Distribution of the total sample according to the demographic characteristics.

(Table-1), The majority of the health care providers joined in the study are of age From the age of thirty and over table 4.1 indicates that the highest percentage is located in age group of (>30) which is (2.36%). According to gender, the sample of the study, which is (74.9%), is male. According to level of qualification all staff which is (40.9%) are institute, and the lowest percentage of nurses with higher degrees (7.1%). Regarding years of experiences in Six years and over years is (73.1 %.)It also showed the highest percentage of medical staff who are married. In addition, those who live in the city.

Items	Rating	F.	%	M.S	S.D	Ass.
Do you know	Correct	157	91.8	1.08	0.275	P

the difference between washing hands and rubbing hands?	Incorrect	14	11.1			
	Total	171	100.0			
Is wearing gloves an alternative to rubbing hands?	Correct	54	31.6	1.68	0.466	M
	Incorrect	117	68.4			
	Total	171	100.0			
Do you routinely rub your hands with alcohol for hand hygiene?	Correct	108	63.2	1.73	0.484	M
	Incorrect	63	36.8			
	Total	171	100.0			
Is it necessary to rub the hands after taking off the paws?	Correct	118	69	1.31	0.464	M
	Incorrect	53	31			
	Total	171	100.0			
Do you ask your senior colleagues or doctors about the process of rubbing hands?	Correct	89	52.0	1.48	0.501	M
	Incorrect	82	48.0			
	Total	171	100.0			
Is there any harm in using surgical paws on your hands?	Correct	69	40.4	1.60	0.492	M
	Incorrect	102	59.6			
	Total	171	100.0			
Are personal protective equipment (gloves, masks, gloves, etc.) always available at your workplace?	Correct	104	60.8	1.39	0.490	M
	Incorrect	67	39.2			
	Total	171	100.0			
	Correct	64	37.4			
	Incorrect	107	62.6			

Table 2. Health Care Providers awareness about Knowledge related to hand hygiene

Do you remember a specific situation that prevented you from applying a hand scrub?	Total	171	100.0	1.63	0.485	M
Have you received training on how to scrub your hands in the last 3 years?	Correct	73	42.7	1.57	0.496	M
	Incorrect	98	57.3			
	Total	171	100.0			
Are you satisfied with your knowledge about hand hygiene?	Correct	130	76.0	1.24	0.428	M
	Incorrect	41	24.0			
	Total	171	100.0			
Do you wash your hands before interacting	Correct	124	72.5	1.27	0.448	M

with the patient?	Incorrect	47	27.5			
	Total	171	100.0			
Do you wash your hands after interacting with a patient?	Correct	164	95.9	1.04	0.199	P
	Incorrect	7	4.1			
	Total	171	100.0			
In your opinion, does the manager of your organization have an impact on improving and enhancing hand-washing procedures?	Correct	78	45.6	1.54	0.500	M
	Incorrect	93	54.4			
	Total	171	100.0			
Is it important to publish hand hygiene posters in your department as a reminder to perform hand hygiene?	Correct	166	97.1	1.03	0.169	P
	Incorrect	5	2.9			
	Total	171	100.0			
To be a good example in performing the process of cleaning hands has an impact on others in this regard.	Correct	159	93.0	1.07	0.256	P
	Incorrect	12	7.0			
	Total	171	100.0			

Table 3.

Do you need a reminder in order to apply the process of cleaning hands?	Correct	65	38.0	1.62	0.487	M
	Incorrect	106	62.0			
	Total	171	100.0			
Is it important for the patient to ask you to clean hands?	Correct	99	57.9	1.42	0.495	M
	Incorrect	72	42.1			
	Total	171	100.0			

Table 4.

Assessment levels (0.0_0.38) poor;(0.39_0.63) moderate;(0.64-1.00) good

Table (2) Appear that the overall health care providers approaches are associated to HH. The study result indicates that the overall evaluation acceptable awareness at mean score (1.39).

Results of table(2) shows that there were a the answers in the questionnaire about realization procedures and information of hand hygiene acceptable cognizance by the health care staff, and showed an unsatisfactory result the staff's difference between washing hands and rubbing hands, and wash hands after interacting with a patient is considered the basic link to reduce and suppress infection and prevent the spread of pathogens within the health institution, so care and awareness must be taken towards, and showed a high score of About the use of personal protective equipment such as gloves, masks, etc. in the workplace.

Discussion

Part I: Discussion of the Demographic Characteristics of the Study Sample, as Shown in Table (1) With respect to age group, the study finding revealed that the majority of study group were 30 years old and above. This finding agrees with that of The knowledge of hand hygiene among the healthcare workers of two teaching hospitals in Mashhad [15] who stated that the participants located in age group of (>30) constitute the majority. Concerning of

gender, the study finding revealed that majority in the study group were male. This finding is supported by [16 [17] also these results are consistent with [18], who showed in their study that most of the staff are males at a rate of 75% and 25% for females. In relation to educational qualification, the study finding displayed that majority of study group medical staff were associate degree graduates (institute). This finding was supported by [19]. Regarding the years of experience, the study finding indicated that the highest percentage in the study concerning years of experiences Six years and over. This finding was congruent with [20]. That who concluded that the years of experience in nursing were 5-7 years.

Part II: Discussion of the Overall Health Care Providers awareness about Knowledge related to hand hygiene, as Show in Tables (2):

In this section, awareness of hand hygiene was examined using a questionnaire according to the guidelines of the World Health Organization. The level of fulfillment of all information was of an average range of approximately at mean score (1.39). The results are consistent with previous findings [21-23]. They also showed poor practices a difference between hand washing and hand rubbing, and the result was similar to other studies as in [24]. Participants also considered the use of alcohol for hand hygiene to be an essential and important factor, which is consistent with [25-27]. They also considered that using gloves can be considered an alternative to rubbing hands because it is easy to use and does not require any effort, and this agrees with [28-29]. This sample demonstrated its ability to promote access to hand hygiene products. This is consistent with the previous study [30].

The evaluation results showed that there were no instructions or educational materials to update their knowledge about hand hygiene. They also seemed unprepared for the presence of future posters for health practices, and the lack of personal protective equipment among individuals in all Al Nasiriyah hospitals and their failure to adhere to it, in addition to the absence of continuous monitoring and evaluation. This is what the study results in principle indicated.

Conclusion

The study indicated that there is fair level of awareness of health staff in all hospitals in Nasiriyah Governorate toward (HH) with Poor hand washing and hygiene after Assessing the patient without any precautionary measures, and showed an unacceptable aspect in the absence of instructions, directives or future strategies regarding (personal protective equipment, sterilization techniques, etc.). There must be a system to improve the performance of health institutions to spread the commitment of health care staff and spread awareness about hand hygiene.

References

1. . D. J. Gould, D. Moralejo, N. Drey, J. H. Chudleigh, and M. Taljaard, "Interventions to improve hand hygiene compliance in patient care," *Cochrane Library*, vol. 2017, no. 9, Sep. 2017, doi: 10.1002/14651858.cd005186.pub4.
2. . M. L. Grayson et al., "Outcomes from the first 2 years of the Australian National Hand Hygiene Initiative," *The Medical Journal of Australia*, vol. 195, no. 10, pp. 615-619, Nov. 2011, doi: 10.5694/mja11.10747.
3. . T. -t. Huang and S. -c. Wu, "Evaluation of a training programme on knowledge and compliance of nurse assistants' hand hygiene in nursing homes," *Journal of Hospital Infection*, vol. 68, no. 2, pp. 164-170, Jan. 2008, doi: 10.1016/j.jhin.2007.11.020.
4. . M. C. Sendall, L. K. McCosker, and K. Halton, "Cleaning Staff's Attitudes about Hand Hygiene in a Metropolitan Hospital in Australia: A Qualitative Study," *International Journal of Environmental Research and Public Health*, vol. 16, no. 6, p. 1067, Mar. 2019, doi: 10.3390/ijerph16061067.
5. . N. Wischnewski, M. Mielke, and C. Wendt. "Healthcare-associated infections in long-term care facilities (HALT) in Frankfurt am Main, Germany, January to March 2011," *PubMed*, Aug. 30, 2012. <https://pubmed.ncbi.nlm.nih.gov/22958607/>.
6. . E. Tacconelli and M. Pezzani, "Public health burden of antimicrobial resistance in Europe.," 2019.
7. . C. Hughes, M. Tunney, and M. C. Bradley, "Infection control strategies for preventing the transmission of meticillin-resistant *Staphylococcus aureus* (MRSA) in nursing homes for older people," *Cochrane Library*, vol. 2013, no. 11, Nov. 2013, doi: 10.1002/14651858.cd006354.pub4.
8. . D. Pittet, "Infection control and quality health care in the new millenium," *American Journal of Infection Control*, vol. 33, no. 5, pp. 258-267, Jun. 2005, doi: 10.1016/j.ajic.2004.11.004.
9. . N. S. Baldwin et al., "Cluster randomised controlled trial of an infection control education and training intervention programme focusing on meticillin-resistant *Staphylococcus aureus* in nursing homes for older people," *Journal of Hospital Infection*, vol. 76, no. 1, pp. 36-41, May 2010, doi: 10.1016/j.jhin.2010.03.006.
10. . M.-L. Ho, W.-H. Seto, L.-C. Wong, and T.-Y. Wong, "Effectiveness of Multifaceted Hand Hygiene Interventions in Long-Term Care Facilities in Hong Kong: A Cluster-Randomized Controlled Trial," *Infection Control and Hospital Epidemiology*, vol. 33, no. 8, pp. 761-767, Jul. 2012, doi: 10.1086/666740.
11. . W. K. Yeung, W. S. W. Tam, and T. W. Wong, "Clustered Randomized Controlled Trial of a Hand Hygiene Intervention Involving Pocket-Sized Containers of Alcohol-Based Hand Rub for the Control of Infections in Long-Term Care Facilities," *Infection Control and Hospital Epidemiology*, vol. 32, no. 1, pp. 67-76, Nov.

- 2010, doi: 10.1086/657636.
12. . J. Hammerschmidt and T. Manser, "Nurses' knowledge, behaviour and compliance concerning hand hygiene in nursing homes: a cross-sectional mixed-methods study," *BMC Health Services Research*, vol. 19, no. 1, Aug. 2019, doi: 10.1186/s12913-019-4347-z.
 13. . A. Huis, L. Schoonhoven, R. Grol, R. Donders, M. Hulscher, and T. Van Achterberg, "Impact of a team and leaders-directed strategy to improve nurses' adherence to hand hygiene guidelines: A cluster randomised trial," *International Journal of Nursing Studies*, vol. 50, no. 4, pp. 464-474, Aug. 2012, doi: 10.1016/j.ijnurstu.2012.08.004.
 14. . E. Alp, A. Ozturk, M. Guven, I. Celik, M. Doganay, and A. Voss, "Importance of structured training programs and good role models in hand hygiene in developing countries," *Journal of Infection and Public Health*, vol. 4, no. 2, pp. 80-90, Jun. 2011, doi: 10.1016/j.jiph.2011.03.001.
 15. . H. Zakeri, F. Ahmadi, E. Rafeemanesh, and L. A. Saleh, "The knowledge of hand hygiene among the healthcare workers of two teaching hospitals in Mashhad," *Electronic Physician*, vol. 9, no. 8, pp. 5159-5165, Aug. 2017, doi: 10.19082/5159.
 16. . M. Desta, T. Ayenew, N. Sitotaw, N. Tegegne, M. Dires, and M. Getie, "Knowledge, practice and associated factors of infection prevention among healthcare workers in Debre Markos referral hospital, Northwest Ethiopia," *BMC Health Services Research*, vol. 18, no. 1, Jun. 2018, doi: 10.1186/s12913-018-3277-5.
 17. . R. D. I. Aledeilah, N. M. A. El-Fetoh, and A. F. Albaker, "Assessment of Knowledge, Attitude and Practice of Hand Hygiene among Health Care Workers in Arar City, Saudi Arabia," *The Egyptian Journal of Hospital Medicine*, vol. 70, no. 3, pp. 491-498, Jan. 2018, doi: 10.12816/0043494.
 18. . H. I. Faris and H. B. Hassan, "Evaluation of Nurses Practices Concerning Sterile Techniques Critical Care Units in Al-Najaf AL-Ashraff City Hospitals," *journal-article*. [Online].
 19. . H. I. Faris and H. B. Hassan, "Evaluation of Nurses Practices Concerning Sterile Techniques Critical Care Units in Al-Najaf AL-Ashraff City Hospitals," *journal-article*. [Online].
 20. . D. Pittet, "Infection control and quality health care in the new millenium," *American Journal of Infection Control*, vol. 33, no. 5, pp. 258-267, Jun. 2005, doi: 10.1016/j.ajic.2004.11.004.
 21. . A. Sharif, A. Arbabisarjou, A. Balouchi, S. Ahmadidarrehshima, and H. H. Kashani, "Knowledge, Attitude, and Performance of Nurses toward Hand Hygiene in Hospitals," *Global Journal of Health Science*, vol. 8, no. 8, p. 57, Dec. 2015, doi: 10.5539/gjhs.v8n8p57.
 22. . KM. Abd Elaziz, IM, Bakr, "Assessment of knowledge, attitude and practice of hand washing among health care workers in Ain Shams University hospitals in Cairo," *PubMed*, Mar. 01, 2009. <https://pubmed.ncbi.nlm.nih.gov/19771756/>
 23. . J. Ahmed et al., "Compliance and Knowledge of Healthcare Workers Regarding Hand Hygiene and Use of Disinfectants: A Study Based in Karachi," *Cureus*, Feb. 2020, doi: 10.7759/cureus.7036.
 24. . V. Maheshwari, N. C. M. Kaore, V. K. Ramnani, S. K. Gupta, A. Borle, and R. Kaushal, "A Study to Assess Knowledge and Attitude Regarding Hand Hygiene amongst Residents and Nursing Staff in a Tertiary Health Care Setting of Bhopal City," *Journal Of Clinical And Diagnostic Research*, Jan. 2014, doi: 10.7860/jcdr/2014/8510.4696.
 25. . World Health Organization, "WHO Guidelines on Hand Hygiene in Health Care: a Summary," 2009. [Online]. Available: <https://www.pediatrics.gob.mx/archivos/burbuja/guia2.pdf>
 26. . Guidelines Review Committee, "WHO guidelines on hand hygiene in health care," Jan. 15, 2009. <https://www.who.int/publications/i/item/9789241597906>
 27. . G. T. Engdaw, M. Gebrehiwot, and Z. Andualem, "Hand hygiene compliance and associated factors among health care providers in Central Gondar zone public primary hospitals, Northwest Ethiopia," *Antimicrobial Resistance and Infection Control*, vol. 8, no. 1, Nov. 2019, doi: 10.1186/s13756-019-0634-z.
 28. . K. Vinodhini and B. A, "Study On Infection Control Practices Among Healthcare Workers In A Speciality Hospital, Chennai," *ResearchGate*, Jun. 2016, [Online].
 29. . H. S. Vikke et al., "Hand hygiene perception and self-reported hand hygiene compliance among emergency medical service providers: a Danish survey," *Scandinavian Journal of Trauma Resuscitation and Emergency Medicine*, vol. 27, no. 1, Feb. 2019, doi: 10.1186/s13049-019-0587-5.
 30. . I. R. Okafor, "Hand hygiene knowledge and practices among healthcare providers in a tertiary hospital, south west, Nigeria," *www.academia.edu*, Jan. 2013, [Online]. Available: https://www.academia.edu/104391743/Hand_hygiene_knowledge_and_practices_among_healthcare_providers_in_a_tertiary_hospital_south_west_Nigeria