

Table Of Content

Journal Cover 2

Author[s] Statement 3

Editorial Team 4

Article information 5

 Check this article update (crossmark) 5

 Check this article impact 5

 Cite this article 5

Title page 6

 Article Title 6

 Author information 6

 Abstract 6

Article content 8

Academia Open



By Universitas Muhammadiyah Sidoarjo

Originality Statement

The author[s] declare that this article is their own work and to the best of their knowledge it contains no materials previously published or written by another person, or substantial proportions of material which have been accepted for the published of any other published materials, except where due acknowledgement is made in the article. Any contribution made to the research by others, with whom author[s] have work, is explicitly acknowledged in the article.

Conflict of Interest Statement

The author[s] declare that this article was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright Statement

Copyright © Author(s). This article is published under the Creative Commons Attribution (CC BY 4.0) licence. Anyone may reproduce, distribute, translate and create derivative works of this article (for both commercial and non-commercial purposes), subject to full attribution to the original publication and authors. The full terms of this licence may be seen at <http://creativecommons.org/licences/by/4.0/legalcode>

EDITORIAL TEAM

Editor in Chief

Mochammad Tanzil Multazam, Universitas Muhammadiyah Sidoarjo, Indonesia

Managing Editor

Bobur Sobirov, Samarkand Institute of Economics and Service, Uzbekistan

Editors

Fika Megawati, Universitas Muhammadiyah Sidoarjo, Indonesia

Mahardika Darmawan Kusuma Wardana, Universitas Muhammadiyah Sidoarjo, Indonesia

Wiwit Wahyu Wijayanti, Universitas Muhammadiyah Sidoarjo, Indonesia

Farkhod Abdurakhmonov, Silk Road International Tourism University, Uzbekistan

Dr. Hindarto, Universitas Muhammadiyah Sidoarjo, Indonesia

Evi Rinata, Universitas Muhammadiyah Sidoarjo, Indonesia

M Faisal Amir, Universitas Muhammadiyah Sidoarjo, Indonesia

Dr. Hana Catur Wahyuni, Universitas Muhammadiyah Sidoarjo, Indonesia

Complete list of editorial team ([link](#))

Complete list of indexing services for this journal ([link](#))

How to submit to this journal ([link](#))

Article information

Check this article update (crossmark)



Check this article impact (*)



Save this article to Mendeley



(*) Time for indexing process is various, depends on indexing database platform

The Efficacy of a Three-Week Low-Calorie Diet in Decreasing Liver Volume Pre-Laparoscopic Sleeve Surgery: A Randomized Clinical Trial

Efektivitas Diet Rendah Kalori Selama Tiga Minggu dalam Mengurangi Volume Hati Pra-Bedah Lengan Laparoskopik: Uji Klinis Secara Acak

Sura Aqeel Sadiq, sura@gmail.com, (0)

M.B.Ch. B, Basrah Health Office /Primary Health Department, Iraq

Sajjad Salim Issa, sajjad.issa@uobasrah.edu.iq, (1)

Community Health Nursing Department, College of Nursing, University of Basrah, Basrah, Iraq, Iraq

Jasim D. Saud, jasim@gmail.com, (0)

M. B. Ch. B, F.A.B.H.S, F.I.B.M.S, I.F.S.O, Consultant surgeon, Basrah Health Office/Basrah Teaching Hospital, Iraq

⁽¹⁾ Corresponding author

Abstract

Background: For obese individuals having significant non-bariatric surgery, better preoperative weight loss techniques are required. Objectives: To identify the efficacy of a three-week low-calorie diet in decreasing liver volume pre-laparoscopic sleeve surgery. Methods: Twenty-five morbidly obese individuals who were scheduled for LSG participated in this interventional trial. Every patient was instructed to adhere to a low-calorie, restrictive diet for three weeks, and their liver span was measured by ultrasonography before and after the regimen. The end measurements were the effects of the preoperative diet on liver span and liver enzymes (ALT and AST). Results: Liver span is significantly reduced as a function of diet. Weight and body mass index showed a favorable correlation with changes in liver size. Liver enzyme levels significantly increased. Conclusion: A low-calorie diet successfully decreased liver size for three weeks before bariatric surgery.

Highlights:

1. Evaluate 3-week low-calorie diet's effect on liver size before surgery.

2. Obese patients followed low-calorie diet; liver span measured pre/post.

3. Diet reduced liver size, correlated with weight, but increased liver enzymes.

Keywords: Sleeve Surgery, Liver Volume, Three weeks, Efficacy

Academia Open

Vol 9 No 1 (2024): June

DOI: 10.21070/acopen.9.2024.10368 . Article type: (Clinical Research)

Published date: 2024-06-28 00:00:00

Introduction

Obesity is the accumulation and abnormal distribution of body fat brought on by many factors, including genetics, a diet heavy in calories and fat, and a lack of physical activity [1].

Obesity has been a major public health concern in both developed and developing regions of the world. Over 650 million of the 1.9 billion overweight persons in the world in 2016 were obese, a number that has tripled since the 1970s. According to projections, if the pace keeps increasing, by 2025, over one-third of adults worldwide will be overweight, and over 1 billion will be obese. By 2030, up to 57.8% of the world's adult population, or 3.3 billion people, will be either overweight or obese [2].

Obesity is defined as a person's body mass index (BMI) of 30 kg/m² or above. This figure is calculated by dividing the square of the height in meters by the body weight in kilos [3].

Coronary artery disease, hyperlipidemia, hypertension, sleep apnea, and non-alcoholic fatty liver disease are all associated with high body mass index [4].

When bariatric surgery is used to treat extreme obesity, it can improve and remit several obesity-related comorbidities, cause long-term weight loss, enhance quality of life, and increase survival [5].

A very low-calorie diet is commonly characterized as consuming 450-800 kcal per day, whereas a low-calorie diet consumes 800- 1500 kcal per day [6,7]. The duration of very low-calorie diets ranges from 2 weeks to 8 weeks, and the consistency ranges from entirely liquid meal replacements in the form of commercially available protein shakes to a combination of liquid meal replacement and food meals [8,9,10].

Methods

The prospective interventional study design has been adopted to evaluate the effectiveness of 3 weeks of a low-calorie diet in a patient awaiting bariatric surgery. The study is conducted at two hospitals: Basrah Teaching Hospital and AL-Sader Teaching Hospital. The study period started from Feb 2023 to the end of June 2023.

Forty men and women who were morbidly obese and candidates for laparoscopic sleeve gastrectomy bariatric surgery made up the study population. Men and women aged 20-55 who met the following inclusion criteria were eligible: (a) had a body mass index > 35 kg/m² and were associated with comorbid conditions like type 2 diabetes mellitus, hypertension, dyslipidemia, and sleep apnea; or (b) had a body mass index >40 kg/m² and had not responded to conservative treatment. The following were the exclusion criteria: (a) BMI less than 35; (b) insulin treatment due to hypoglycemia risk; (c) coagulation issues; and (d) serious organ or systemic disease.

Patients undergoing surgery in the surgical wards of Al-Sader Teaching Hospital and Basrah Teaching Hospital provided the data. The patients underwent a physical examination both before and after the low-calorie diet was implemented, had their weight recorded, had a blood sample taken for a liver function test, and had an upper abdominal ultrasound before and after the diet.

For three weeks, a low-calorie diet consisting of 800-1200 kcal per day is prescribed. The liver is evaluated by abdominal ultrasonography and blood is drawn to measure the liver function tests ALT and AST.

A general surgeon provided dietary guidance and monitoring, and the participants were instructed to maintain the same level of physical activity during the three-week diet, which included 800-1200 kcal [11].

Result and Discussion

Result

When classified by sex, Table 1 indicates that neither the age nor the levels of BMI, liver span, ALT, and AST showed any statistically significant differences.

Sex	Age	BMI	BMI	Liver	Liver	ALT	ALT	AST	AST	
		before diet	after diet	span before diet	span after diet	before diet	after diet	before diet	after diet	
Male	N	8	8	8	8	8	8	8	8	
	Mean±S	37.75±9.	43.86±4.	42.55±4.	15.5±1.1	13.59±1.	31.5±16.	32.12±1	25.75±1	26.54±1
	D	25	56	22	9	23	08	6.02	4.34	4.28

Academia Open

Vol 9 No 1 (2024): June

DOI: 10.21070/acopen.9.2024.10368 . Article type: (Clinical Research)

	Median (Min.-Max.)	-----	-----	-----	-----	13 (12-15.7)	28.5 (13-60)	28.8 (14.1-60.5)	22.5 (12-55)	23.1 (13-55.6)
Female	N	17	17	17	17	17	17	17	17	17
	Mean±SD	39.29±9.22	43.73±3.79	42.19±3.48	15.35±1.62	13.79±1.65	26.73±9.62	27.48±9.56	22.88±8.85	23.68±8.86
	Median (Min.-Max.)	-----	-----	-----	-----	13 (11.6-17)	26 (7-50)	27 (8-51)	22 (8-46)	22.1 (9-47)
P-value		0.700	0.939	0.826	0.821	0.929	0.466	0.485	0.930	0.907

Table 1. Comparison of the levels of the study variables categorized

** Mann-Whitney U

		BMI before diet	BMI after diet	Liver span before diet	Liver span after diet	ALT before diet	ALT after diet	AST before diet	AST after diet
Age	R	-.143-	-.137-	.184	.351	.084	.090	.153	.147
	Sig.	.496	.514	.378	.085	.691	.669	.466	.482
	N	25	25	25	25	25	25	25	25
BMI before diet	R		.990**	.235	.001	.401*	.403*	.398*	.399*
	Sig.		.000	.258	.998	.047	.046	.049	.048
	N		25	25	25	25	25	25	25
BMI after diet	R			.243	.009	.409*	.410*	.399*	.401*
	Sig.			.242	.968	.043	.042	.048	.047
	N			25	25	25	25	25	25
Liver span before the diet	R				.871**	.246	.250	.297	.298
	Sig.				.000	.235	.228	.150	.148
	N				25	25	25	25	25
Liver span after diet	R					.019	.020	.055	.057
	Sig.					.927	.925	.792	.787
	N					25	25	25	25
ALT before diet	R						.999**	.944**	.942**
	Sig.						.000	.000	.000
	N						25	25	25
ALT after diet	R							.945**	.944**
	Sig.							.000	.000
	N							25	25
AST before diet	R								1.000**
	Sig.								.000
	N								25

Table 2. Spearman's and Pearson's correlations between the variables under investigation

A substantial statistical difference between the BMI before and after diet regimens was discovered when the change in BMI was examined for significance. Following the diet, the BMI dramatically dropped.

	Mean±SD	N	Paired Mean±SD	P-value
BMI before diet	43.77±3.96	25	1.46±0.61	0.0001
BMI after diet	42.31±3.65	25		

Table 3. Tests of paired BMI to determine the change post the dietary regimen

	Liver span before the diet	Liver span after diet	ALT before diet	ALT after diet	AST before diet	AST after diet
N	25	25	25	25	25	25

Mean±SD	15.4±1.47	13.72±1.51	28.26±11.93	28.97±11.86	23.8±10.68	24.59±10.66
Median (Min.-Max.)	16 (13-18)	13 (11.6-17)	27 (7-60)	27 (8-60.5)	22 (8-55)	22.1 (9-55.6)

Table 4. The study variables' mean and median values

		N	Mean Rank	P-value
Liver span after diet - Liver span before diet	Negative Ranks	25a	13.00	0.0001
	Positive Ranks	0b	0.00	
	Ties	0c		
	Total	25		
ALT after diet - ALT before diet	Negative Ranks	1d	2.50	0.0001
	Positive Ranks	24e	13.44	
	Ties	0f		
	Total	25		
AST after diet - AST before diet	Negative Ranks	0g	0.00	0.0001
	Positive Ranks	25h	13.00	
	Ties	0i		
	Total	25		

Table 5. shows the changes in paired liver span, ALT, and AST tests following the food course.

* Wilcoxon Signed Ranks Test

- a. Liver span after diet < Liver span before diet
- b. Liver span after diet > Liver span before diet
- c. Liver span after diet = Liver span before diet
- d. ALT after diet < ALT before diet
- e. ALT after diet > ALT before diet
- f. ALT after diet = ALT before diet
- g. AST after diet < AST before diet
- h. AST after diet > AST before diet
- i. AST after diet = AST before die

Discussion

Regarding the results of the study, most of the study sample ages were between 30 to 39 years old. The results of the present study agree with other studies [12-16] which reveal that most of the study ages were between 30 to 39 years old.

Regarding the results of the study, most of the study sample were female. The results of the current study agree with other studies [17-23] which reveal that most of the study sample were female.

According to the study, liver size was significantly reduced (11%) by following a low-calorie diet for three weeks before laparoscopic sleeve gastrectomy, similar to many studies [24-28].

Both the starting BMI and liver size, as well as the weight and BMI after the diet, were positively connected with the reduction in liver size. These results agreed with many studies [29-35].

The BMI decreased by 1 to 1.5 kg/m² Preoperative weight loss is recommended by various studies to lower the risk of postoperative complications. The results of this study agreed with a study [36] which revealed the majority of patients showed a considerable rise in their liver enzymes (ALT and AST). The degree of cytolysis in those following an LCD diet may help to explain this.

According to the study by [37], this rise is temporary during the first few weeks on LCD, and related metrics did not alter significantly after 30 weeks. Since ultrasound is a quick, simple, and affordable method that works well in underdeveloped nations, we decided to use it to test the liver span. It has been demonstrated that liver size and

texture can be assessed using ultrasonography [38].

The target population of our study was extremely uniform in terms of the level of obesity and dietary composition, which is one of its strengths. It may be questioned in the future if a universal LCD will be advantageous for all patients. Depending on the objectives being set by a multidisciplinary team, customized preoperative diets might be preferable. Patients with varying BMIs or comorbidities may have varied objectives, such as stabilizing glucose levels or reducing liver volume. New research examining the impact of LCDs in various study populations is justified.

Conclusion

This study shows that an LCD can effectively lower the weight and volume of the liver. It is advised that an LCD lasts for three weeks and delivers 800-1500 kcal per day. According to earlier research using a VLCD, an LCD may even be useful for reducing liver volume. An LCD should therefore be used since it can prevent needless, severe dietary restriction and its aftereffects (such as adverse effects).

References

1. . A. M. Tiryag and H. H. Atiyah, "Nurses' Knowledge Toward Obesity in Al-Basra City," *Annals of the Romanian Society for Cell Biology*, vol. 25, no. 3, pp. 4667-4673, May 2021.
2. . S. S. Issa, S. M. Ibrahim, and A. A. Al-Mussawi, "A Study on Abdominal Obesity at Basra University Staffs," *Clinical Medicine Research*, vol. 6, no. 3, pp. 69-73, 2017.
3. . F. A. Jassim, Z. T. Maki, and S. S. Issa, "Prevalence of Overweight and Obesity Among Primary School Children in Basrah City Center," *HIV Nursing*, vol. 23, no. 2, pp. 152-157, Jan. 2023.
4. . E. H. Rahi, Z. M. Al-Hejaj, and A. M. Tiryag, "Nurses' Knowledge of Nonalcoholic Fatty Liver Disease: A Cross-Sectional Study," *Academia Open*, vol. 9, no. 1, pp. 10-21070, Jun. 2024.
5. . A. M. Tiryag and H. H. Atiyah, "Nurses' Knowledge Toward Bariatric Surgery at Surgical Wards at Teaching Hospitals in Al-Basra City," *Indian Journal of Forensic Medicine & Toxicology*, vol. 15, no. 3, pp. 5152-5159, Jun. 2021.
6. . E. Delbridge and J. Proietto, "State of the Science: VLED (Very Low Energy Diet) for Obesity," *Asia Pacific Journal of Clinical Nutrition*, vol. 15, suppl. 1, pp. 49-54, 2006.
7. . M. Holderbaum, D. S. Casagrande, S. Sussenbach, et al., "Effects of Very-Low-Calorie Diets on Liver Size and Weight Loss in the Preoperative Period of Bariatric Surgery: A Systematic Review," *Surgery for Obesity and Related Diseases*, vol. 14, no. 2, pp. 237-244, 2018.
8. . J. Ahmed and S. S. Issa, "Association of Abdominal Obesity with Dietary Habits and Demographic Variables Among Students at Southern Technical University," *University of Thi-Qar Journal of Science*, vol. 10, no. 1 (SI), Apr. 2023.
9. . S. S. Issa, H. H. Abdul-Ra'aoof, F. A. Jassim, M. M. Al-Khalissi, A. A. Munther, A. A. Alwan, and M. B. Abd, "A Study About Obesity Among Preschool Children at Al-Zubair District," *HIV Nursing*, vol. 22, no. 2, pp. 3403-3407, Nov. 2022.
10. . A. K. Dekan, J. T. Ahmed, and S. S. Issa, "Association of Overweight and Obesity with Dietary Habits and Some Socio-Demographic Variables Among Students in Southern Technical University," *International Journal of Health Sciences*, vol. 3, pp. 6856-6870, 2024.
11. . M. A. Abdullah, F. J. Kadhum, and S. S. Issa, "Assessment of Liver Involvement in Patients with Type 2 Diabetes Mellitus in Basrah City, Iraq, Using FibroScan and Correlation with Risk Factors," *Cureus*, vol. 16, no. 7, Jul. 2024.
12. . Z. K. Dhahi, S. S. Issa, and L. A. Hameed, "A Study on Pregnant Women's Satisfaction with Primary Health Care Services in Basra," *International Journal of Research in Humanities, Arts and Literature*, vol. 3, no. 1, pp. 7-19, Jan. 2015.
13. . S. S. Issa, K. J. Madwah, and H. S. Al Mosawi, "Evaluation of Nurses' Knowledge in Management of Premature Babies in Neonatal Units," *American Journal of Nursing*, vol. 6, no. 5, pp. 291-295, 2018.
14. . S. M. Ebrahim and S. S. Issa, "Workplace Violence Against Nursing Staff Working in Emergency Departments at General Hospitals in Basra City," *Indian Journal of Public Health Research & Development*, vol. 9, no. 6, pp. 239-244, 2018.
15. . S. M. Ebrahim and S. S. Issa, "Satisfaction with Nursing Care Among Patients Attending Oncology Center in Basra City, Iraq," *Journal of Environmental Science and Engineering*, vol. 4, pp. 241-248, 2015.
16. . H. H. Abdul-Ra'aoof, S. B. Dawood, F. A. Jassim, S. K. Jassim, S. S. Issa, M. A. Akber, and M. A. Atiyah, "Moderate Proficiency in Suture Techniques Among Nurses: A Cross-Sectional Study," *Journal of Nursing Education and Practice*, vol. 14, no. 2, pp. 50-58, 2024.
17. . H. H. Abdul-Ra'aoof, M. A. Akber, F. A. Jassim, A. M. Tiryag, S. S. Issa, M. A. Atiyah, J. A. Mezail, and I. S. Hassan, "The Psychological Impact of Violence on Emergency Department and Intensive Care Unit Nurses: A Cross-Sectional Study," *Research Journal of Trauma and Disability Studies*, vol. 3, no. 4, pp. 228-233, Apr. 2024.
18. . H. M. Sabty, S. B. Dawood, and A. M. Tiryag, "Nurses' Knowledge and Practices on Influenza Vaccination for Pregnant Women," *Jurnal Kebidanan Midwiferia*, vol. 10, no. 2, pp. 50-59, Oct. 2024.

Academia Open

Vol 9 No 1 (2024): June

DOI: 10.21070/acopen.9.2024.10368 . Article type: (Clinical Research)

19. . A. M. Tiryag, "Revitalizing Hearts: The Transformative Impact of Pacemaker Therapy on Cardiac Conduction Disorders," *Academia Open*, vol. 9, no. 1, pp. 10-21070, Jun. 2024.
20. . I. H. Zainel, H. H. Abdul-Ra'aoof, and A. M. Tiryag, "Mothers' Knowledge and Attitudes Towards Their Children with Neonatal Jaundice: A Cross-Sectional Study," *Health Education and Health Promotion*, vol. 10, no. 3, pp. 565-570, Jul. 2022.