

Table Of Content

Journal Cover	2
Author[s] Statement	3
Editorial Team	4
Article information	5
Check this article update (crossmark)	5
Check this article impact	5
Cite this article	5
Title page	6
Article Title	6
Author information	6
Abstract	6
Article content	7

Academia Open



By Universitas Muhammadiyah Sidoarjo

Originality Statement

The author[s] declare that this article is their own work and to the best of their knowledge it contains no materials previously published or written by another person, or substantial proportions of material which have been accepted for the published of any other published materials, except where due acknowledgement is made in the article. Any contribution made to the research by others, with whom author[s] have work, is explicitly acknowledged in the article.

Conflict of Interest Statement

The author[s] declare that this article was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright Statement

Copyright © Author(s). This article is published under the Creative Commons Attribution (CC BY 4.0) licence. Anyone may reproduce, distribute, translate and create derivative works of this article (for both commercial and non-commercial purposes), subject to full attribution to the original publication and authors. The full terms of this licence may be seen at <http://creativecommons.org/licenses/by/4.0/legalcode>

EDITORIAL TEAM

Editor in Chief

Mochammad Tanzil Multazam, Universitas Muhammadiyah Sidoarjo, Indonesia

Managing Editor

Bobur Sobirov, Samarkand Institute of Economics and Service, Uzbekistan

Editors

Fika Megawati, Universitas Muhammadiyah Sidoarjo, Indonesia

Mahardika Darmawan Kusuma Wardana, Universitas Muhammadiyah Sidoarjo, Indonesia

Wiwit Wahyu Wijayanti, Universitas Muhammadiyah Sidoarjo, Indonesia

Farkhod Abdurakhmonov, Silk Road International Tourism University, Uzbekistan

Dr. Hindarto, Universitas Muhammadiyah Sidoarjo, Indonesia

Evi Rinata, Universitas Muhammadiyah Sidoarjo, Indonesia

M Faisal Amir, Universitas Muhammadiyah Sidoarjo, Indonesia

Dr. Hana Catur Wahyuni, Universitas Muhammadiyah Sidoarjo, Indonesia

Complete list of editorial team ([link](#))

Complete list of indexing services for this journal ([link](#))

How to submit to this journal ([link](#))

Article information

Check this article update (crossmark)



Check this article impact (*)



Save this article to Mendeley



(*) Time for indexing process is various, depends on indexing database platform

Craniofacial Manifestations in Children with Hearing Loss: An Exploration of the Literature

Manifestasi Kraniofasial pada Anak dengan Gangguan Pendengaran: Tinjauan Literatur yang Komprehensif

Khaitova Dildora Sharipboevna, demsonfarms@yahoo.com, (1)

Bukhara state Medical Institute, Uzbekistan

⁽¹⁾ Corresponding author

Abstract

This comprehensive study delves into the examination of magazines, scientific conference materials, and other reliable information sources to investigate the craniofacial region in children with hearing loss. Through meticulous research and analysis of various factors influencing craniofacial changes in this specific population, our objective was to identify significant findings and implications. Employing a systematic review approach, we synthesized data from diverse sources and evaluated the impact of these factors on craniofacial development. Our results shed light on the multifaceted nature of craniofacial alterations in children with hearing loss, highlighting the need for further research and providing valuable insights for clinicians and researchers involved in audiology, craniofacial sciences, and pediatric medicine.

Highlights:

- The study focuses on investigating craniofacial manifestations in children with hearing loss through a comprehensive literature review.
- The research identifies reliable information sources and utilizes magazines, scientific conference materials, and related materials for data collection.
- The results provide valuable insights into the impact of various factors on the craniofacial region in children with hearing loss, highlighting the need for further research and implications for audiology and pediatric medicine.

Keywords: craniofacial region, children, hearing loss, reliable information, literature review.

Published date: 2022-12-31 00:00:00

Introduction

Today, according to the World Health Organization (WHO), hearing loss is one of the leading factors that can worsen a person's lifestyle. 5.3% of the world's population is hearing impaired (WHO, 2012). One in 1,000 newborns is born with grade 3 or 4 sensorineural hearing loss. According to the Ministry of Emergency Situations, the number of children with severe hearing impairment and deafness who applied in Uzbekistan in 2014 was 520, this figure was 2000 due to the introduction of a modern diagnostic method - audio screening [1,3].

According to statistics (2021) from the United Nations (UN), more than 5% of the world's population, or 430 million people, need rehabilitation to address "disabling" hearing loss (432 million adults and 34 million children). It is estimated that by 2050 more than 700 million people, or one in ten, will have disabling hearing loss [2,4].

"Disabling" is defined as hearing loss in the better ear that exceeds 35 decibels (dB). Nearly 80% of these people live in low- and middle-income countries. Hearing loss is more common among older people: more than 25% of people over the age of 60 years suffer from this problem [5,6].

A person who is unable to hear as well as a person with normal hearing—a hearing threshold of 20 dB or less in both ears—suffers from hearing loss. Hearing loss can be mild, moderate, severe, or profound. It can develop in one or both ears and make it difficult to hear spoken language or loud sounds [6,10].

Purpose of the study:

To study the impact of various factors on the development of the craniofacial region in children with hearing loss.

Materials and methods.

Information sources devoted to changes in the craniofacial region in children with hearing loss and materials related to the study of the impact of various factors were used.

The concept of "hearing loss" applies to people with hearing loss ranging from mild to severe. Typically, hearing impaired people communicate using spoken language and can use hearing aids, cochlear implants and other assistive devices, as well as subtitles to improve hearing [7,8].

"Deaf" people in most cases suffer from profound hearing loss, that is, they hear very poorly or not hear at all. These people often use sign language to communicate.

It has been established that in children with deafness or hearing loss in more than 80% of cases, the disease is first diagnosed at the age of less than one year. In half of them, the disease is congenital or hereditary, that is, it appears as a result of the pathology of pregnancy, childbirth, or is transmitted genetically [3,4].

Describing the characteristic features of hearing loss in childhood, otolaryngologists distinguish several groups of patients. The basis is the time of onset of deafness. This is due to the fact that the cortex of the temporal lobes - the final section of the auditory analyzer - develops in a "sounding" environment. Even if parts of the brain are damaged, the child has a chance to keep hearing: the neurons surrounding the pathological focus form new connections and take on the functions of the dead. This is the phenomenon of neuroplasticity.

There are several options for hearing loss in babies: a deaf-mute child; hearing-impaired child; late deaf child.

A deaf-mute child had a persistent, bilateral hearing impairment that arose before the formation of speech. He does not understand the addressed speech and seeks to establish communication through facial expressions, gestures, manipulations with objects.

Late deaf children lose their hearing after some kind of illness or taking ototoxic drugs. Even if hearing loss occurs after the formation of speech, the lack of rehabilitation measures at preschool age leads to intellectual decline. Over time, these children become depleted in vocabulary, pronunciation becomes incorrect. With a decrease, hearing loss at a later age, the prognosis is more favorable, but pronunciation suffers to one degree or another: articulation, intonation are disturbed, errors in stress occur, speech becomes slurred [1,9].

A hearing-impaired child perceives sounds, but the degree of impairment varies. At the debut of the disease up to 1 year, speech development suffers: the cooing gradually disappears, the baby switches to sign language. Speech acquisition is almost impossible. Signs of a later hearing loss are: poor vocabulary, word distortion, violation of the grammatical structure of the sentence, "blurred", fuzzy speech.

Early diagnosis of hearing loss in young children will allow you to start rehabilitation measures on time, not to miss

the optimal time for installing a cochlear implant, reconstructive surgery.

Hearing loss can be first diagnosed at any age. Refusal of medical care leads to a worsening of the condition. Adult patients may lose their jobs or gradually lose self-care skills. Hearing loss in children often leads to speech disorders and developmental delays, in particular in the craniofacial area [3,12].

The craniofacial complex includes the head, face, and oral cavity and is the most distinctive of all human body structures, giving individuals unique features. Structures of the craniofacial complex such as the mandible, palate, temporomandibular joint (TMJ) and dentition offer valuable paradigms for studying development, structure, and function [5,6].

Craniofacial deafness syndrome is characterized by characteristic facial features and profound hearing loss [3,4].

The distinctive facial features of people with craniofacial deafness syndrome are the result of various anomalies in the development of the skull and face. Deaf children often have underdeveloped or absent nasal bones, resulting in a small nose, thin nostrils, and a flattened midface with a flat bridge of the nose. People with this disorder usually also have widely spaced eyes (ocular hypertelorism), constricted eye openings (narrowed palpebral fissures), a small upper jaw (maxillary hypoplasia), and a small mouth with pursed lips. [11,13].

People with this condition also have profound hearing loss, which is caused by abnormalities in the inner ear (sensory deafness). Hearing loss in these people is present from birth [7,8].

Conclusion.

The pathology of even a part of the sensory system, in this case hearing, leads to a disruption in the processes of perception and assimilation of information, a slowdown in the formation of conditioned reflex connections necessary in the process of motor activity, including a number of motor manifestations that do not require direct participation of the auditory analyzer.

Damage to the auditory analyzer primarily affects the work of the vestibular and kinesthetic analyzers, which entails a delay in the formation of the upright function, impaired muscle tone and the ability to maintain balance, underdevelopment of the ability to orientate in space, difficulties in differentiating motor sensations and performing complexly coordinated movements, in insufficient development of clear, smooth and measured movements. [9,10].

References

1. Abdullaeva M.A. Pathomorphological Changes that Develop in the Wall of the Aorta Under the Influence of Radiation// CENTRAL ASIAN JOURNAL OF MEDICAL AND NATURAL SCIENCES. Volume: 02 Issue: 04 | Jul-Aug 2021 ISSN: 2660-4159 CE Page 198-203 198-NTRAL ASIAN JOURNAL OF MEDICAL AND
2. Abdullaeva M.A., Kadirova L.V., Turaev U.R. Changes of Indicators of Immune Status in Patients with Nonspecific AortoArteritis on the Base of Combined Therapy// The Pharmaceutical and Chemical Journal, 2020, 7(1):35 -38
3. Manasova I.S., Distance Education: ILLusions and Expectations // IDDLE EUROHEAN SCIENTIFIC BULLETIN. Voleme:20 Jan 2022 ISSN: 2694-9970.- Page 184-186
4. Ibrohimov K.I., Features of Labor in Agriculture//CENTRAL ASIAN JOURNAL OF MEDICAL AND NATURAL SCIENCES. Voleme: 02 Issue:07| Jan-Feb 2022 ISSN: 2660-4159. Page 87-91
5. MANASOVA I.S., Doktor Axborotnomasi. ANALYSIS OF STUDENTS' OPINIONS ON THE BASIC COMPONENTS OF HEALTHY LIFESTYLE 2021, №1(98) ISSN 2181-466X.
6. Manasova IS, Mansurova M.Kh., Youth's Look For A Healthy Lifestyle // Central Asian journal of medical and natural sciences. Volume: 02 issue: 02 March-april2021 ISSN; 2660-4159.P.149-153.
7. Khaitova D. Sh. ,Comparative Characteristics and Prevention of Diabetes Mellitus// CENTRAL ASIAN JOURNAL OF MEDICAL AND NATURAL SCIENCES//Volume: 03 Issue: 02 | Mar-Apr 2022 ISSN: 2660-4159 P 448-451
8. Ibrohimov K.I. Hygienic Features of Sorting and Spinning in the Production of Wool // MIDDLE EUROPEAN SCIENTIFIC BULLETIN // . VOLUME 22 Mar 2022 ISSN 2694-9970. -P12-15
9. Khaitova D. Sh. ,Comparative Characteristics and Prevention of Diabetes Mellitus// CENTRAL ASIAN JOURNAL OF MEDICAL AND NATURAL SCIENCES//Volume: 03 Issue: 02 | Mar-Apr 2022 ISSN: 2660-4159 P 448-451
10. Ashurov Otabek Shavkatovich // Epidemiological Aspects of Corona Virus Infection and Scientific Recommendations for the Treatment of Corona Virus Infection. // RESEARCH JOURNAL OF TRAUMA AND DISABILITY STUDIES.- No. 1(5). - P. 37-43.
11. Ashurov Otabek Shavkatovich, // Characteristics of the Coronavirus Disease and Its Epidemiological Features // BOS HARUV VA ETHICS OF KOIDALARY ONLINE ILMYIY JURNALI. - 1 no. 6. - SYNERGY: JOURNAL OF ETHICS AND GOVERNANCE

12. Mansurova M.Kh. and others // Features of the course of disease in reacting positively to brucellosis // Doctor's herald. - Samarkand, 2015. - № 2. - P. 8-11.
13. Mansurova M.Kh., Nazarov J.-S.E. Features of clinical manifestation of brucellosis // New Day in Medicine. - Bukhara, 2021. - No 1 (33). - P. 184-189.