## Academia Open Vol 9 No 2 (2024): December

Vol 9 No 2 (2024): December DOI: 10.21070/acopen.9.2024.8121 . Article type: (Science)

# **Table Of Content**

Journal Cover	2
Author[s] Statement	3
Editorial Team	4
Article information	5
Check this article update (crossmark)	5
Check this article impact	5
Cite this article	5
Title page	6
Article Title	6
Author information	6
Abstract	6
Article content	7



By Universitas Muhammadiyah Sidoarjo

ISSN 2714-7444 (online), https://acopen.umsida.ac.id, published by Universitas Muhammadiyah Sidoarjo Copyright © Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC

> BY). 2/18

Vol 9 No 2 (2024): December DOI: 10.21070/acopen.9.2024.8121 . Article type: (Science)

#### **Originality Statement**

The author[s] declare that this article is their own work and to the best of their knowledge it contains no materials previously published or written by another person, or substantial proportions of material which have been accepted for the published of any other published materials, except where due acknowledgement is made in the article. Any contribution made to the research by others, with whom author[s] have work, is explicitly acknowledged in the article.

## **Conflict of Interest Statement**

The author[s] declare that this article was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

## **Copyright Statement**

Copyright © Author(s). This article is published under the Creative Commons Attribution (CC BY 4.0) licence. Anyone may reproduce, distribute, translate and create derivative works of this article (for both commercial and non-commercial purposes), subject to full attribution to the original publication and authors. The full terms of this licence may be seen at <a href="http://creativecommons.org/licences/by/4.0/legalcode">http://creativecommons.org/licences/by/4.0/legalcode</a>

Vol 9 No 2 (2024): December DOI: 10.21070/acopen.9.2024.8121 . Article type: (Science)

## **EDITORIAL TEAM**

## **Editor in Chief**

Mochammad Tanzil Multazam, Universitas Muhammadiyah Sidoarjo, Indonesia

## **Managing Editor**

Bobur Sobirov, Samarkand Institute of Economics and Service, Uzbekistan

### Editors

Fika Megawati, Universitas Muhammadiyah Sidoarjo, Indonesia Mahardika Darmawan Kusuma Wardana, Universitas Muhammadiyah Sidoarjo, Indonesia Wiwit Wahyu Wijayanti, Universitas Muhammadiyah Sidoarjo, Indonesia Farkhod Abdurakhmonov, Silk Road International Tourism University, Uzbekistan Dr. Hindarto, Universitas Muhammadiyah Sidoarjo, Indonesia Evi Rinata, Universitas Muhammadiyah Sidoarjo, Indonesia M Faisal Amir, Universitas Muhammadiyah Sidoarjo, Indonesia Dr. Hana Catur Wahyuni, Universitas Muhammadiyah Sidoarjo, Indonesia

Complete list of editorial team (link) Complete list of indexing services for this journal (link) How to submit to this journal (link)

Vol 9 No 2 (2024): December DOI: 10.21070/acopen.9.2024.8121 . Article type: (Science)

# **Article information**

## Check this article update (crossmark)



## Check this article impact <sup>(\*)</sup>



## Save this article to Mendeley



 $^{(\ast)}$  Time for indexing process is various, depends on indexing database platform

Vol 9 No 2 (2024): December DOI: 10.21070/acopen.9.2024.8121 . Article type: (Science)

# Novel Synthesis of Thiopyrimidine Compounds from Dibenzo-18-Crown-6 and Evaluation of Antibacterial Activity

#### Zena Abdali , zenaabdaliabdali@gmail.com, (1)

Department of Pharmaceutical Chemistry, College of Pharmacy, University of Basrah, Iraq

#### Raheem Jameel Mahesein, mahesein@gmail.com, (0)

Department of Pharmaceutical Chemistry, College of Pharmacy, University of Basrah, Iraq

<sup>(1)</sup> Corresponding author

#### Abstract

In this study, we present a novel synthesis approach for the production of thiopyrimidine compounds, includingP-OH-Thiopyrimidine,P-NO2-Thiopyrimidine, andP-Cl2-Thiopyrimidine, from the precursor compound dibenzo-18-crown-6 (DB-18-CE-6). These thiopyrimidine compounds have shown significant potential for various applications, particularly as antibacterial agents. The synthesis involved the reaction of DB-18-CE-6-derived alpha-beta unsaturated ketones with thiourea in a basic medium, resulting in successful cyclization. The synthesized thiopyrimidine compounds were evaluated for their antibacterial activity against common bacterial strains, including Streptococcus, Staphylococcus, and Proteus, and compared to the antibiotic Azithromycin. The results demonstrated that the thiopyrimidine compounds (Z, Z1, Z2, Z3) exhibited excellent antibacterial activity, surpassing Azithromycin, and highlighting their potential as effective alternatives to combat bacterial infections. This research opens new avenues for the development of antibacterial agents and emphasizes the significance of thiopyrimidine compounds in addressing bacterial resistance issues.

#### **Highlights** :

- Novel synthesis of thiopyrimidine compounds from DB-18-CE-6.
- Evaluation of antibacterial activity against Streptococcus, Staphylococcus, and Proteus.
- Potential of thiopyrimidine compounds as effective antibacterial agents.

**Keywords :** Thiopyrimidine compounds, Antibacterial activity, DB-18-CE-6, Novel synthesis, Bacterial resistance.

Published date: 2023-10-11 00:00:00

# Introduction

Crown ethers are mecrocyclic polyethars that are wall known for the nancovalent ion banding capabilities that they possess. They were first reported by Pedersen [1-3], . The term "classical crown ethers" refers to a class of macrocyclic polyethers containing anywhere from three to twinty oxygon atoms, each separated from the next by twe or more carben atoms [4]. The numbir and typa of atoms in the polyethar ring, the class nime "crown" that cames from thair molecular shape to "crewn" a matal ien upen coerdination, and the number of oxygon atems in the polyethar ring. In general, common names are preferred over systematic names. Take, for instance, the compound dibenzo-18-crown-6 [5]. These molecules' binding selectivity for a variety of metal ions is determined by the size of their crown rings; as a result, the 18-crown-6 molecule has a strong affinity for potassium, whereas the 15-crown-5 molecule prefers sodium cations. In order to provide an explanation for this phenomenon, it has been discovered that high complex stability is linked to a larger penatration of the metel catian into the polyather civity [6, 7].

Crewn ethar macrecycles are knewn in all ring sizes, ranging from 9 to at least 60, which has led to a large deal of structural diversity (exceeding 10,000 examples) [8]. As a result, crown ethers have been the subject of a significant amount of research from a variety of perspectives, such as senser applications, biological medel systims [9],. Because crown ether derivatives work in a manner that is analogous to that of naturally occurring ionophores (such as gramicidin), researchers have been able to utilize them to investigate a variety of biological processes. In particular, researchers have found that crown ether derivatives have the ability to create channels and transport ions through lipid membranes [9]. Crown ethers can be functionalized by a variety of chemical processes due to their high solubility. Thus, numerous crown ether derivatives have been manufactured. Conformational flexibility is one of the distinguishing characteristics of crown ethers.

In recent years, dibenzo-18-crown-6 has garnered attention in the field of organic synthesis, particularly for its potontial a versatile buelding bleck fir the synthesis of functionalized compounds. Its unique structural features make it a suitable starting material for various transformations, such as acylation, alkylation, and cyclization reactions [10].

In the present work the cyclization of thiopyrimidine of dibenzo-18-crown-6, was accomplished by the reaction of alfa beta unsaturated ketone of DB-18-CE-6 who synthesis by dissolve (0.444g, 0. 1 mmole) of acytal-DB-18-CE-6 in 5ml of absolute ethanol then added (3-5 drops) of (60%KOH in ethanol) slowly with constant stirring follow added (0.106g, 0.2mmole) benzyldehyde which dissolved in absolute ethanol. The reaction mixture was kept in refrigerator overnight. The product was filtered and washed with cold water

until the filtered were neutral to litmus, the product was dried and recrystilized to get pale yellow color product [1]. As in the reaction who attended previously in first paper of synthesis, characterization, and preliminary of new antimicrobial activity of alfa beta unsaturated ketone of dibenzo-18-crown-6. Follow reaction with thiourea ,the mixture of alfa-beta unsaturated ketone (0.31g 0.001mmole) and thiourea (0.152g 0.002mmole) dissolve in absolute ethanol 5ml, with 5 drops of (40% KOH in ethanol) wes edded slewly with constent strring. The reection mexture allewed to reflex in a watar beth .. The ppt. obtained was filtered, washed with water, and dried. The product was recrystallized using dichloromethane, dried overnigh [2]. The goal is to synthesis cyclization thiopyrimidine of DB-18- CE-6 compounds to explore the potential antibacterial of these compounds.

In this experiment we made compared between the thiopyrimidine compounds of DB-18-CE-6 and control azithromycin antibiotic which is macrolide antibiotic.

Macrolide antibiotics are known for their wide range of efficacy. They are effective against Gram-positive and Gramnegative bacteria in equal measure. Although the action spectrum of macrolides may vary very slightly from one another azithromycin was one of the drugs that was used the most at the onset of the SARS-CoV2 epidemic. Because it encourages the development of drug-resistant bacterial strains. Regarding azithromycin, the main mechanism for increasing bacterial resistance to this antibiotic is changing the target of the active ingredient on the ribosome [11, 12].

# **Material and methods**

Alfa beta unsaturated ketone of DB-18-CE-6compounds, thiourea, KOH. The progress of reaction was monitored by TLC (Silica gel 60 F254). IR spectra were recorded on Perkin-Elmer spectrometer. The  $^{1}$ HNMR , $^{13}$ CNMR with Chloroform/d/DMSO-d6 as solventand TMS as internal standard. Melting points were determined in an open capillary tube.

#### Synthesis of thiopyrimidine of DB-18-CE-6 (Z ) $% \left( {Z_{\rm{B}}} \right)$

The mixture of alfa-beta unsaturated ketone of DB-18-CE-6 (0.31g 0.1mmole) and thiourea (0.152g 0.2mmole) dissolve in absolute ethanol 5ml, with 5 drops of (40% KOH ) in watar bith In between TLG wes monitored chock the complation of the reection., let to dried overnight. fully charecterized by elemantal anilysis (FTt-IR,<sup>1</sup>Hh-NMR,

BY). 7/18

and <sup>13</sup>Cc-NMR).



4-phenyl-6-(14-(6-phenyl-2-thioxo-2H-1<sup>2</sup>-pyrimidin-4-yl)-6,7,9,10,17,18,20,21-octahydrodibenzo[b.k][1,4,7,10,13,16]hexaoxacyclooctadecin-2-yi)pyrimidin-2(1H)-thione

#### Figure 1. Scheme 1 Thiopyrimidine dibenzo-18-crown-6 synthesis

#### Synthesis of P-OH-thiopyrimidine of dibenzo-18-crown- 6(Z 1)

The mixture of alfa-beta unsaturated ketone of DB-18-CE-6 (0.32g 0.1mmole) and thiourea (0.152g 0.2mmole) dissolve in absolute ethanol 5ml, with 5 drops of (40% KOH in ethanol) wes addid slewly weth constint stiring. The rection mixtur allowed to reflex in a water bath In between TLG was menitored to chieck the complition of the rection. aftr compltion of the rection, let to dried overnight. fully characterized by elemental analysis (FTt-IRr, <sup>1</sup>Hh-NMRr, and <sup>13</sup>Cc-NMR).

#### Synthesis of p-NO2-thiopyrimidine of dibenzo-18-crown- 6(Z 2)

The mixture of alfa-beta unsaturated ketone of DB-18-CE-6 (0.355g 0.1mmole) and thiourea (0.152g 0.2mmole) dissolve in absolute ethanol 5ml, with 5 drops of (40% KOH in ethanol) let to dried overnight. fully characterized by elemental analysis (FTt-IRr,  $^{1}$ Hh-NMRr, and  $^{13}$ Cc-NMR).

#### Synthesis of p-cl2-thiopyrimidine of dibenzo-18-crown- 6(Z 3)

The mixture of alfa-beta unsaturated ketone of DB-18-CE-6 (0.344g, 0.1 mmole) and thiourea (0.152g, 0.2 mmole) dissolve in absolute ethanol 5ml, with 5 drops of (40% KOH in ethanol) lit to dried overnight. fully characterized by elemental analysis (FT-IR, <sup>1</sup>H-NMR, and <sup>13</sup>C-NMR).

#### Antibacterial activity

The antibacterial potential of the prepared Samples (Thiopyrimidine compunds of dibenzo-18- crown-6Z, Z1, Z2, Z3, Azithromycin) was investigated against Gram's negative and Gram's positive bacterial strains using agar well diffusion assay [13]. Into the bored wells, different concentrations (50, 150, 200, 250, 400 ppm) of the Samples (Z, Z1, Z2, Z3, Azithromycin) were used. The cultured plates containing the Samples (Z, Z1, Z2, Z3, Azithromycin, and the test organisms were incubated overnight at 37°C before measuring and recording the average the zones of inhibition diameter [14].

# **Result and Discussion**

Crown ethers can be functionalized by a variety of chemical processes due to their high solubility. Thus, numerous crown ether derivatives have been manufactured. Conformational flexibility is one of the distinguishing characteristics of crown ethers [8]. The focus on the cyclization of alfa beta unsaturated ketones of dibenzo-18-crown-6 by reaction with thiourea into thiopyrimidine of dibenzo-18-crown-6, The goal is to explore the potential antibacterial of these compounds. By utilizing dibenzo-18-crown-6 as a precursor, it is expected to obtain thiopyrimidine of dibenzo-18-crown-6 compounds with enhanced biological activity and potentially novel mechanisms of action [5, 8, 16].

In this experiment cyclization of thiopyrimidine of dibenzo-18-crown-6, was accomplished by the reaction of alfa beta unsaturated ketone of dibenzo-18-crown-6 who synthesis by dissolve (0.222mg, 0.5mmole) of acytaldibenzo-18-crown-6-ether in 5ml of absolute ethanol then added (3-5 drops) of (60%KOH in ethanol) slowly with good stirring follow by added (0.106g, 1mmole) of benzyldehyde dissolved it in ethanol absolute was added to the mixture slowly with for 2hr. Then added ice crush from deionized water then filter and let dry at R.T follow reaction with thiourea.in basic medium.

#### Copyright @ Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC

Clasien-Schmidt condensation:



#### Figure 2. Scheme 2 General mechanism of thiopyrimidine dibenzo-18-crown-6 synthesis

All the newly produced derivative compounds Z, Z1, Z2, Z3, have FT-IR spectra that share similarities in certain fingerprint-like bands and other bands. The essential functional group vibration bands are presented in Table (1 and 2) and the corresponding compound IR spectra are displayed in Figures (1) to (4).

All the predicted bands for normal Compound, as well as the bonded of add groups are present in all FT-IR spectra of derivatives.

The carboxylic groups bond of O-H present in Z3, performing a robust and broad extending band vibration in the range of 3367-3450 cm<sup>-1</sup>. Furthermore, the N-H bond can be assigned a medium band in the range of 3275-3488 cm<sup>-1</sup>. The compounds Z-Z3 of IR spectra show a band at a range of 3064-3172 cm<sup>-1</sup> because of aromatic straining C-H, as well as 3 powerful bands to moderate ones at the range of 997.2-736.81 cm<sup>-1</sup> because of the bending of C-H bond aromatic. A weak band was observed at the range of 2949-2835 cm<sup>-1</sup> due to asymmetrical straining of aliphatic C-H bands, whereas a moderate band emerged at the range of 1398-1325 cm<sup>-1</sup> for the bending of aliphatic C-H bond. Aromatic (C=C) asymmetrical and symmetrical stretching could be associated with two prominent bands that popped up in the 1544-1455 cm-1 and 1597-1415 cm-1 ranges, and between. Further to that, the compounds and Z-Z3 of FT-IR spectra illustrate a strong band that is attributed to (C=N) at the range of 1724-1610 cm<sup>-1</sup> and assigned to (C-O) at the range of 1085- 1051 Cm<sup>-1</sup> of all compound.

	O-HStr etching	N-HStr etching	C-HStre tching( Aromati c)	C-HStre tching( aliphati c)	C=N	C=CStr etching	C-HBend hat	ling(Alip tic)	C-O-CSt retchin g	C-OStre tching	C-HBen ding(Ou t plane)	C-O-Hb ending
Z	-	3381.21 3275.13	3172.90	2927.94 2877.79	1610.56	1415.75	-1504.48	1325.10	1126.43 1215.15 1251.8	1060.85 1085.92	943.197 36.81	626.874 89.92
Z1	-	3439.08 3419.79	3062.96	2949.16 2927.94 2877.79	1724.36 1616.35	1597.06 145	1506.41 4.33	1398.39 1340.53 1328.95	1253.73 1217.08 1128.36	1058.92	997.295 6.69943 .19779. 24774.5 2	599.86
Z2	-	3421.72	3064.89	2949.16 2926.01 2883.58	0	1595.13 145	1510.26 4.33	1330.88	1290.38 1255.66 1232.51 1132.21	1060.85 1051.20	997.293 5.48779 .24740. 67	597.934 97.63
Z3	3367-3 448	3448-3 367	3064.89	2949.16 2926.01 2883.58	0	1597.06 145	1510.26 4.33	1330.88	1290.38 1255.66 1230.58 1132.21	1058.92	997.293 5.48779 .24742. 59	597.934 97.63

ISSN 2714-7444 (online), https://acopen.umsida.ac.id, published by Universitas Muhammadiyah Sidoarjo

Copyright © Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY).

Vol 9 No 2 (2024): December DOI: 10.21070/acopen.9.2024.8121 . Article type: (Science)

#### Table 1. The data and vibrational mode description of FTIR spectra of Z-Z3

The antibacterial potential of the prepared samples (Z, Z1, Z2, Z3) and Azithromycin were investigated against Gram's negative and Gram's positive bacterial strains using agar well diffusion assay. 6 mm-diameter wells were bored on the agar plates using of a sterile tip. Into the bored wells, different concentrations of the Samples (Z, Z1, Z2, Z3, Azithromycin) were used. The cultured plates containing the samples (Z, Z1, Z2, Z3, Azithromycin) were incubated overnight at 37°C before measuring and recording the average the zones of inhibition diameter [14], will show in the tables below.

The rise in antibiotic resistance that can be attributed to a number of different sources has led researchers to explore for novel chemicals that are effective against infections that are resistant to several drugs. In this experiment. Thiopyrimidine compounds displayed a diversity of pharmacological properties, and its derivatives possess a high degree of structural variance. As a result, there is a critical demand for the development of novel antibacterial medication candidates [17]. Antibiotic resistance is a rising problem, reducing the effectiveness of macrolides, the first-line treatment for many diseases. Staphylococci frequently exhibit multiple molecular determinants of resistance to macrolides. Research into new macrolide antibiotics could help in the fight against drug-resistant bacteria, hence it's crucial to understand the processes of macrolide resistance. The comparative analysis with azithromycin indicated the potential of thiopyrimidine compounds as an effective alternative in resist bacterial infection .<sup>(11, 18)</sup> As in the scheme below show the bacterial inhibition for the thiopyrimidine compounds with different concentration against Streptococcus, Staphylococcus and Proteus as we see in concentration 50pmm,100ppm,150ppm,200ppm,400ppm [10]. The bacterial inhibition for thiopyrimidine compound (Z) in (400ppm) against Streptococcushas inhibition zone (22mm) larger than Azithromycin in (400ppm) have inhibition zone(14mm) that is mean the effect of thiopyrimidine compound have a good antibacterial activity against streptococcus, While inhibition zone of compound (Z) in (400ppm) against Stapholococcus is (12mm) for Azithromycin the inhibition zone is (14mm) And the inhibition zone for compound (Z) against

*Proteus*is (11mm), for Azithromycin is (16mm).

# Conclusions

Evaluation of activity Conduct comprehensive antibacterial activity assays using a range of bacterial strains. Determine the minimum inhibitory concentrations (MICs) of thiopyrimidine compounds against various pathogens to assess their potency and efficacy. Comparative studies with known antibiotics like Azithromycin can provide valuable insights into their relative effectiveness.

Table (3) Explain the antibacterial activity of thiopyrimidine compounds of dibenzo-18-crown-6 Z=Thiopyrimidine of DB-18-CE-6, Z1=P-OH-thiopyrimidine of DB-18-CE-6, Z2=P-NO2- thiopyrimidine, Z3=P-CL-thiopyrimidine

sample		А	В	С	D	Е	F
Streptococc us	Z	6	15	17	18	20	22
	Z1	6	16	17.5	18.5	19.5	20.5
	Z2	6	15	16	17	18	19
	Z3	6	16	18	20	22	24
Azithro	omycin	6	8	9	10	13	14
Staphylococ cus	Z	6	6	7	8	10	12
	Z1	6	6	6.5	7	9	10
	Z2	6	7	8	9	12	13
	Z3	6	6.5	8	9	12	13
Azithro	omycin	6	7	9	10	12	14
Proteus	Z	6	6	7	9	10	11
	Z1	6	6	7	10	11	12
	Z2	6	9	10	11	12	13
	Z3	6	6	7	9	12	13
Azithro	omycin	6	12	13	14	15	16

 Table 2. Antibacterial analysis (Zone of inhibition)

Copyright © Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC

Academia Open Vol 9 No 2 (2024): December DOI: 10.21070/acopen.9.2024.8121 . Article type: (Science)



Figure 3. The FTIR Spectra of compound Z

11/18

Academia Open Vol 9 No 2 (2024): December DOI: 10.21070/acopen.9.2024.8121 . Article type: (Science)



Figure 4. The FTIR Spectra of compound Z1

Academia Open Vol 9 No 2 (2024): December DOI: 10.21070/acopen.9.2024.8121 . Article type: (Science)



Figure 5. The FTIR Spectra of compound Z2

13/18

Academia Open Vol 9 No 2 (2024): December DOI: 10.21070/acopen.9.2024.8121 . Article type: (Science)



Figure 6. The FTIR spectra of compound Z3

Academia Open Vol 9 No 2 (2024): December DOI: 10.21070/acopen.9.2024.8121 . Article type: (Science)



Figure 7. The 1H NMR spectrum of c ompound Z2

Academia Open Vol 9 No 2 (2024): December DOI: 10.21070/acopen.9.2024.8121 . Article type: (Science)



Figure 8. The 1H NMR spectrum of c ompound Z1

Copyright © Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY).

16/18

Academia Open Vol 9 No 2 (2024): December DOI: 10.21070/acopen.9.2024.8121 . Article type: (Science)



Figure 9. The 1H NMR spectrum of c ompound Z3

# **References**

- 1. D. Septianingtyas, N. Zafira, Z. Zulhipri, F. Kurniadewi, H. Dianhar, Eds., "Green synthesis of chalcones derivatives," in AIP Conference Proceedings, 2021.
- 2. B. M. Sahoo, M. Rajeswari, P. Jnyanaranjan, S. Binayani, "Green expedient synthesis of pyrimidine derivatives via chalcones and evaluation of their anthelmintic activity," Indian Journal of Pharmaceutical Education and Research, vol. 51, no. 4S, pp. S700-S6, 2017.
- R. Izatt, "Synthetic multidentate macrocyclic compounds," Elsevier, 2012.
   G. W. Gokel, W. M. Leevy, M. E. Weber, "Crown ethers: sensors for ions and molecular scaffolds for materials and biological models," Chemical reviews, vol. 104, no. 5, pp. 2723-2750, 2004.
- 5. M. Febles, S. Montalvão, G. D. Crespín, M. Norte, J. M. Padrón, P. Tammela, et al., "Synthesis and biological evaluation of crown ether acyl derivatives," Bioorganic & Medicinal Chemistry Letters, vol. 26, no. 22, pp. 5591-5593, 2016.
- 6. M. K. Marwan, J. A. Layla, "Synthesis, Characterization and Biological studies of new crown ether derivatives," 2009.

Copyright @ Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC

- 7. W. M. Leevy, G. M. Donato, R. Ferdani, W. E. Goldman, P. H. Schlesinger, G. W. Gokel, "Synthetic Hydraphile Channels of Appropriate Length Kill Escherichia coli," Journal of the American Chemical Society, vol. 124, no. 31, pp. 9022-9023, 2002.
- 8. M. Kralj, L. Tušek-Božić, L. Frkanec, "Biomedical potentials of crown ethers: prospective antitumor agents," ChemMedChem: Chemistry Enabling Drug Discovery, vol. 3, no. 10, pp. 1478-1492, 2008.
- 9. H. A. Younis, "Design, Synthesis, Characterization and biological studies of New mono, dipeptide Prodrugs, and their Bioactivity" 2023.
- J. P. Rufino, L. G. V. Sousa, A. E. L. Rodrigues, A. L. S. Falcão, I. C. de Melo Lima, L. A. de Miranda Cavalcanti, et al., "Bacterial resistance to azithromycin: causes, effects, and the fight against COVID-19," Research, Society and Development, vol. 11, no. 6, pp. e27711629198-e27711629198, 2022.
- 11. H. H. Bahjat, R. A. Ismail, G. M. Sulaiman, M. S. Jabir, "Magnetic field-assisted laser ablation of titanium dioxide nanoparticles in water for anti-bacterial applications," Journal of Inorganic and Organometallic Polymers and Materials, pp. 3649-3656, 2021.
- 12. M. A. Jihad, F. T. Noori, M. S. Jabir, S. Albukhaty, F. A. AlMalki, A. A. Alyamani, "Polyethylene glycol functionalized graphene oxide nanoparticles loaded with nigella sativa extract: a smart antibacterial therapeutic drug delivery system," Molecules, vol. 26, no. 11, pp. 3067, 2021.
- M. S. Jabir, T. M. Rashid, U. M. Nayef, S. Albukhaty, F. A. AlMalki, J. Albaqami, et al., "Inhibition of Staphylococcus aureus α-hemolysin production using nanocurcumin capped Au@ ZnO nanocomposite," Bioinorganic Chemistry and Applications, vol. 2022, 2022.
- 14. J. Rebek Jr, J. Trend, R. Wattley, S. Chakravorti, "Allosteric effects in organic chemistry. Site-specific binding," Journal of the American Chemical Society, vol. 101, no. 15, pp. 4333-4337, 1979.
- 15. M. Xu, P. Wu, F. Shen, J. Ji, K. Rakesh, "Chalcone derivatives and their antibacterial activities: Current development," Bioorganic Chemistry, vol. 91, pp. 103133, 2019.
- 16. A. T. Feßler, Y. Wang, C. Wu, S. Schwarz, "Mobile macrolide resistance genes in staphylococci," Plasmid, vol. 99, pp. 2-10, 2018.