

Table Of Content

Journal Cover	2
Author[s] Statement	3
Editorial Team	4
Article information	5
Check this article update (crossmark)	5
Check this article impact	5
Cite this article	5
Title page	6
Article Title	6
Author information	6
Abstract	6
Article content	8

Academia Open



By Universitas Muhammadiyah Sidoarjo

Originality Statement

The author[s] declare that this article is their own work and to the best of their knowledge it contains no materials previously published or written by another person, or substantial proportions of material which have been accepted for the published of any other published materials, except where due acknowledgement is made in the article. Any contribution made to the research by others, with whom author[s] have work, is explicitly acknowledged in the article.

Conflict of Interest Statement

The author[s] declare that this article was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright Statement

Copyright © Author(s). This article is published under the Creative Commons Attribution (CC BY 4.0) licence. Anyone may reproduce, distribute, translate and create derivative works of this article (for both commercial and non-commercial purposes), subject to full attribution to the original publication and authors. The full terms of this licence may be seen at <http://creativecommons.org/licences/by/4.0/legalcode>

EDITORIAL TEAM

Editor in Chief

Mochammad Tanzil Multazam, Universitas Muhammadiyah Sidoarjo, Indonesia

Managing Editor

Bobur Sobirov, Samarkand Institute of Economics and Service, Uzbekistan

Editors

Fika Megawati, Universitas Muhammadiyah Sidoarjo, Indonesia

Mahardika Darmawan Kusuma Wardana, Universitas Muhammadiyah Sidoarjo, Indonesia

Wiwit Wahyu Wijayanti, Universitas Muhammadiyah Sidoarjo, Indonesia

Farkhod Abdurakhmonov, Silk Road International Tourism University, Uzbekistan

Dr. Hindarto, Universitas Muhammadiyah Sidoarjo, Indonesia

Evi Rinata, Universitas Muhammadiyah Sidoarjo, Indonesia

M Faisal Amir, Universitas Muhammadiyah Sidoarjo, Indonesia

Dr. Hana Catur Wahyuni, Universitas Muhammadiyah Sidoarjo, Indonesia

Complete list of editorial team ([link](#))

Complete list of indexing services for this journal ([link](#))

How to submit to this journal ([link](#))

Article information

Check this article update (crossmark)



Check this article impact (*)



Save this article to Mendeley



(*) Time for indexing process is various, depends on indexing database platform

Unmasking Hidden Threats Global Spread of MBL Resistance Exposed

Membuka Kedok Ancaman Tersembunyi Penyebaran Global Resistensi MBL yang Terekspos

Ali Hussain Anaid Taaban, 1hly35508@gmail.com, (1)

Department of Pathological Analysis, Faculty of Science, Dhi Qar University, Iraq

Muntadhar Shakir Neama Ali, muntadhar9796@gmail.com, (0)

Department of Pathological Analysis, Faculty of Science, Dhi Qar University, Iraq

Sura Haidar Mutashar Azgair Azgair, Ssry60233@gmail.com, (0)

Department of Pathological Analysis, Faculty of Science, Dhi Qar University, Iraq

Fatima Muzahim Qasim, fatimamuzahim3131f@gmail.com, (0)

Department of Pathological Analyzes, College of Science, University of Sumer, Iraq

⁽¹⁾ Corresponding author

Abstract

This study aims to establish a routine monitoring system for MBL enzymes to provide timely data to healthcare professionals and policy makers, enabling informed decision making on antibiotic use and resistance management. Using a combination of molecular biology techniques and data analysis, we monitor MBL activity in various institutional settings. The increasing prevalence of multidrug-resistant (MDR) bacteria is a significant threat to public health globally. Metallo-beta-lactamase (MBL), an enzyme that confers resistance to a wide range of beta-lactam antibiotics, is particularly concerning due to its ability to spread rapidly in healthcare and community settings. Despite the importance of this issue, systematic monitoring and understanding of MBL remains inadequate. Our findings reveal a significant, previously unreported presence of MBLs, underscoring the urgent need for targeted antibiotic stewardship programs. The implications of this study emphasize the importance of integrating enzyme monitoring into standard healthcare practices to reduce the spread of MDR bacteria.

Highlights:

- Regular Monitoring: Essential for tracking MBL enzyme prevalence and guiding antibiotic use.
- Advanced Techniques: Molecular biology methods enhance MBL detection and analysis.
- Policy Integration: Crucial for implementing enzyme monitoring in healthcare to combat MDR bacteria spread.

Keywords: MBL Enzymes, Antibiotic Resistance, Healthcare Monitoring, Molecular Biology, Stewardship Programs

Academia Open

Vol 9 No 2 (2024): December

DOI: 10.21070/acopen.9.2024.8994 . Article type: (Microbiology)

Published date: 2024-05-17 00:00:00

Introduction

Comprehension their method of action requires a comprehension of (Figure 1). Its structure is identical to that of acyl-D-alanyl-D-alanine, the usual substrate required for the production of the linear glycopeptide found in bacterial cell walls. The B-lactam binds to the enzyme, inhibiting transpeptidation and the production of cell walls [1].

Antibiotic β -lactam structure is shown in Figure (1). A variety of side chains influence an agent's pharmacological characteristics, degree of activity, spectrum, and resistance to B-lactamases. B. P-lactam rings; C. Thiazolidine rings; C'. Dihydrothiazine rings; D. B-lactamases' sites of action; E. amidase sites of action [2], [3].

Comprehending how they work is contingent upon comprehending what is shown in (Figure 1). The structure of β -lactam is similar to that of acyl-D-alanyl-D-alanine, which is the normal substrate needed for the production of linear glycopeptide present in bacterial cell walls. B-lactam binds to enzyme thus inhibiting transpeptidation and the production of cell wall— leading to their death [4].




Figure 1. Structure of β -lactam antibiotics

Antibiotic β -lactam's structure is represented by Figure (1). An agent's pharmacologic characteristics are affected by a variety of side chains influencing activity, spectrum resistance to B-lactamases because these different sites include P-lactam rings; Thiazolidine rings; Dihydrothiazine rings; B-lactamase action sites amidase action sites [5], [6].

Structure of β -lactam antibiotics. a. Different side chains determine degree of activity, spectrum, pharmacologic properties, resistance to β -lactamases; b. β -lactam ring; c. thiazolidine ring; c'.dihydrothiazine ring; d. site of action of β -lactamases; e. site of action of amidase [7].

1. Principles of Antibiotic Resistance

Four primary mechanisms induce bacterial drug resistance, as shown in Table (1). (1) Drugs inactivated by enzymes generated by bacteria, such as β -lactamases, which break the B-lactam ring of cephalosporins and penicillins, rendering them inert [8]. (2) Bacteria create altered targets that are less affected by the drug (for example, a mutated 30S ribosomal subunit protein may lead to streptomycin resistance, and a methylation 23 srRNA can lead to erythromycin resistance) [9]. (3) Drugs are not able to reach an effective intracellular concentration when bacteria lower their permeability (e.g., alterations in porins might diminish the quantity of Penicillin entering the bacterium) [10]. (4) Applying a "efflux" pump, often referred to as a "multidrug-resistance pump". Bacteria actively export therapeutics [11], [12].

2. Lactam Antibiotics Action

Growing bacteria are effectively targeted by all B-lactam drugs due to the specific inhibition of bacterial cell wall formation [13]. While these antibiotics act broadly within the spectrum of their activity, peptidoglycan synthesis suppression is recognized as their major effect because this macromolecule represents an important structural component for most bacteria [14]. Composed of alternating N-acetylglucosamine (NAG) and N-acetylmuramic acid (NAM) molecules, the linear chains are further cross-linked via peptide bridges which result from enzymatic activity during synthesis [15], [16]. When penicillin-based antibiotics interfere with transpeptidases or any other serine protease that constitutes peptidoglycan assembly machinery, there is no room left for doubt about how they do it: in addition to breaking cross-links between peptide stems coming from different chains and hence weakening

cell walls down to bursting point penicillins bind covalently at active site serines on PBPs which normally participate in cross-linking reactions but cannot continue after such binding takes place [17]. Developing bacteria die as a consequence of compromised integrity [18].




Figure (2): Mechanism of action of β - Lactam Antibiotics .

Figure 2. Mechanism of Action of B-Lactam Antibiotics

Bacterial resistance to B-lactam antibiotics is often caused by the creation of B-lactamases, which bind to and hydrolyze them [19]. These enzymes are intended to preferentially detect and hydrolyze quaternary B-lactam rings, resulting in inactive products that can no longer inhibit transpeptidases or TPs [20].

Escherichia coli was determined to have the original B-lactamase, which hydrolyzed penicillin [21]. Because B-lactamases are extensively dispersed and encoded on mobile genetic components, their proliferation and transmission have increased [22]. As a consequence, it is common to encounter bacteria with up to eight distinct forms of B-lactamases, each of which specifically inhibits a family of antibiotics known as beta-lactamases. With over 4300 distinct kinds of enzymes, it is vital to have a reliable and understandable nomenclature to differentiate between them [23].

3. Bactamases Classification

The Bush-Jacoby classification system and the Ambler classification system (AMB) are the two most often used categorization methods for B-lactamases [24]. Bush-Jacoby-Medeiros divided B-lactamases into three groups and sixteen subgroups according to their hydrolytic and inhibitory characteristics [25]. The newest functional categorization has three categories: Cephalosporinases (category C) comprise group 1.

Describe group 3 (class B) and group 2 (class A and D) metallo- β -lactamases as broad-spectrum serine carbenemases that are resistant to inhibitors [26].

Ambler used a fundamental amino acid sequence homology classification approach to divide B-lactamases into four groups (A, B, C, and D). The active sites of class B enzymes include zinc residues, while class A, C, and D enzymes have serine residues [27]. Table 2 shows two categorization algorithms [28].

4. Metallo B-lactamase

Sabath and Abraham discovered metalloprotease lactamases (MBLS) in 1966, which were later identified as class B lactamases in 1980 [29]. Without zinc ions, these enzymes cannot act. Except for monoaminases, almost all metallo- β -lactamases are multiple metalloenzymes capable of hydrolyzing other metallo- β -lactamases. Commercially available metallo- β -lactamase inhibitors including sulbactam, tazobactam, and clavulanic acid do not inhibit them [30]. Furthermore, they are not inhibited by NXL-104 (avibactam sodium hydrate), a class A and C B-lactamase inhibitor [31]. These bacteria are commonly spread by mobile genetic elements such as transposons, plasmids, and

endoviruses, which may infect and disseminate to a range of bacteria, including Enterobacteriaceae and Acinetobacter spp. [32]. The best-known acquired MBLs are IMP- and VIM-type enzymes, which were identified in the early 1990s. Other forms of acquired MBL enzymes discovered include SPM, GIM, SIM, KHM, NDM, AIM, DIM, SMB, TMB, and FIM [33].

Revealed that the most commonly employed and therapeutically helpful class B enzymes are New Delhi MBL (NDM), iminotropase (IMP), and verona integrin-encoded MBL (VIM) [34].

5. Mechanism of Action of B-lactamase

TEM-1 has been the best researched regarding how β -lactamases hydrolyze β -lactams. TEM-1 β -lactamase hydrolyzes β -lactams by first cleaving their amide bonds in two steps [35].

The positively charged enzyme residue at serine B lactamases then attracts the negatively charged carboxylate group of the B-lactam antibiotic to the active site— where the beta-lactam positions itself correctly and forms significant hydrogen bonds with the enzyme. The residues that make up this interaction in the active site are often referred to as oxyanion pore or electrophilic core. The next step is acylation [36].

Methods

Sample collection and processing: Fifty urine samples from UTI patients were taken between April 1 and May 15, 2023, when the patients were registered at Al-Hussein Hospital in Nasiriyah Province. Ten milliliters of clean-catch midstream urine were transferred into sterile containers first thing in the morning [37]. Each specimen was brought into the lab and left there for an hour; if not, it was refrigerated at 4 °C until processing. A standard urine test was performed on each sample to check for the presence of bacteria, white blood cells, and other impurities. In order to isolate bacteria, urine samples were streaked on agar plates (MacConkey, Blood, and Mannitol Salt). The lab's Vitek equipment was utilized to detect the findings. of the same healthcare institution [38].

1. Analyzing Metallo-Beta-Lactamases (MBLs) Phenotypically

Phenotypic characterization of MBL was done using CDDT. Test isolates were streaked on MH agar and dried for three to five minutes to produce turf cultures with 0.5 McFarland opacity standard. Agar plates had imipenem and imipenem/EDTA combination discs (10/750) overlaid on them (Bioanalysis/Turkey) [39]. The inhibition zones of imipenem and imipenem-EDTA disks were measured after overnight incubation at 37°C — indicating positive MBL based on the paper test results, showing that the inhibitory zone of IMP-EDTA paper increased by approximately 7 mm [40].

Results and Discussion

The distribution of bacterial species was the same across the two patient groups with and without bacteriuria, but no multiple isolates were found [41]. The common pathogenicity of E. coli — in patients with symptomatic urinary tract infection — includes pyelonephritis, cystitis and asymptomatic bacteriuria; risk factors for infection include recent sexual intercourse, diaphragm-spermicide use and a history of recurrent infections [42]. The host immune response plus characteristics of bacteria contribute to determining which patient will develop disease due to uropathogenic E. coli (UPEC)..

Hemolysin is a cytotoxic chemical that helps these organisms invade tissue and is often produced by them. P pili are a specific type of pilus that bind to the P blood group antigen and are produced by strains that cause pyelonephritis and express the K antigen [43]. In terms of MBL results, Pseudomonas aeruginosa was the lowest producer, while Klebsiella pneumoniae and E. coli were the highest producers in the study. 80% of MBLs were generated from a total of 35 isolates. With the recent acquisition of genetic constructs, metallo-B-lactamases have emerged as very potent resistance determinants in drug-resistant Gram-negative bacteria, posing a serious risk to human health [44]. Our results support those of [45], who stated that E. coli produced the most MBL. In contrast, [46] found no MBL-positive E. coli in Zanjan city, Iran.

Conclusion

The awareness of MBL acts as the first indicator that antibiotics must be utilized appropriately to limit the spread of MDR bacterial strains inside these hospitals and communities. A new technique of alerting care providers and policy makers about the nature and development of this sort of resistance is to continually monitor MBL enzymes in these surroundings.

References

1. F. M. Abbas, "Metallo-Beta-Lactamases: A Review," *Annals of R.S.C.*, vol. 25, no. 5, pp. 1308, 2021. Available: <https://www.researchgate.net/publication/351286854>
2. S. B. Almasaudi, "Acinetobacter spp. as Nosocomial Pathogens: Epidemiology and Resistance Features," *Saudi J. Biol. Sci.*, vol. 25, no. 3, pp. 586-596, 2018.
3. N. T. Antunes and J. F. Fisher, "Acquired Class D Beta-Lactamases," *Antibiotics (Basel)*, vol. 3, pp. 398-434, 2014. <https://doi.org/10.3390/antibiotics3030398>
4. Z. M. E. Al-lamy, "Detection of ESBLs in Klebsiella spp. Isolated from Clinical and Environmental Samples at Thi-Qar Province," Master Thesis, Thi-Qar University, Iraq, 2016.
5. L. Balsalobre, A. Blanco, and T. Alarcón, "Beta-Lactams," in *Antibiotic Drug Resistance*, J-L. Capelo-Martinez and G. Igrejas, Eds., John Wiley & Sons Inc., pp. 183, 2020.
6. E. Bettioli and S. Harbarth, "Development of New Antibiotics: Taking Off Finally?" *Swiss Med Wkly*, vol. 145, w14167, 2015.
7. E. R. Bevan, A. M. Jones, and P. M. Hawkey, "Global Epidemiology of CTX-M Beta-Lactamases: Temporal and Geographical Shifts in Genotype," *J. Antimicrob. Chemother.*, vol. 72, no. 8, pp. 2145-2155, 2017.
8. K. Bush and M. J. Macielag, "New Beta-Lactam Antibiotics and Beta-Lactamase Inhibitors," *Expert Opin Ther Pat*, vol. 20, pp. 1277-1293, 2010.
9. K. Bush and G. A. Jacoby, "Updated Functional Classification of Beta-Lactamases," *Antimicrob. Agents Chemother.*, vol. 54, pp. 969-976, 2010.
10. I. L. Camargo, H. M. Neoh, L. Cui, and K. Hiramatsu, "Serial Daptomycin Selection Generates Daptomycin-Nonsusceptible Staphylococcus Aureus Strains with a Heterogeneous Vancomycin-Intermediate Phenotype," *Antimicrob. Agents Chemother.*, vol. 52, pp. 4289-4299, 2008.
11. K. C. Carroll et al., "Jawetz, Melnick and Adelbergs Medical Microbiology," 27th ed., McGraw-Hill Education, USA, 2016.
12. M. Castanheira, P. J. Simner, and P. A. Bradford, "Extended-Spectrum Beta-Lactamases: An Update on Their Characteristics, Epidemiology and Detection," *JAC Antimicrob Resist.*, vol. 2, no. 3, 2021. <https://doi.org/10.1093/jacamr/dlab092>
13. E. Chika et al., "Phenotypic Detection of Metallo-Beta-Lactamase and AmpC Enzymes Among Abattoir Isolates of Escherichia coli and Klebsiella Species in Abakaliki, Nigeria," *Trends in Medicine*, vol. 18, no. 1, 2018.
14. M. Dehshiri et al., "The Frequency of Klebsiella Pneumonia Encoding Genes for CTX-M, TEM-1, and SHV-1 Extended-Spectrum Beta-Lactamases Enzymes Isolated from Urinary Tract Infection," *Ann. Clin. Microbiol. Antimicrob.*, 2018.
15. M. V. Edelstein et al., "Spread of Extensively Resistant VIM-2-Positive ST235 Pseudomonas Aeruginosa in Belarus, Kazakhstan, and Russia: A Longitudinal Epidemiological and Clinical Study," *Lancet Infect Dis*, 2013.
16. A. El Salabi et al., "Genetic and Biochemical Characterization of a Novel Metallo-Beta-Lactamase, TMB-1, from an Achromobacter Xylooxidans Strain Isolated in Tripoli, Libya," *Antimicrob. Agents Chemother.*, 2012.
17. R. W. Finberg and R. Guharoy, "Monobactams," in *Clinical Use of Anti-Infective Agents*, Springer, New York, 2012.
18. M. L. Grayson, "Kucers' The Use of Antibiotics: A Clinical Review of Antibacterial, Antifungal, Antiparasitic, and Antiviral Drugs," Hodder Arnold, London, 2010.
19. A. Hamprecht et al., "Detection of the Carbapenemase GIM-1 in Enterobacter Cloacae in Germany," *J Antimicrob Chemother.*, 2013.
20. D. P. Kateete et al., "Carbapenem Resistant Pseudomonas Aeruginosa and Acinetobacter Baumannii at Mulago Hospital in Kampala, Uganda (2007-2009)," 2016.
21. D. T. King, S. Sobhanifar, and N. C. J. Strynadka, "The Mechanisms of Resistance to Beta-Lactam Antibiotics," in *Handbook of Antimicrobial Resistance*, Springer: New York, NY USA, 2017.
22. B. Kocsis and D. Szabó, "Antibiotic Resistance Mechanisms in Enterobacteriaceae," in *Microbial Pathogens and Strategies for Combating Them: Science, Technology and Education*, A Méndez-Vilas, Ed., Formatex Research Center, Spain, 2013.
23. W. Levinson, "Antimicrobial Drugs: Mechanism of Action," in *Review of Medical Microbiology and Immunology*, McGraw-Hill Education, 2016.
24. L. K. Logan and R. A. Weinstein, "The Epidemiology of Carbapenem-Resistant Enterobacteriaceae: The Impact and Evolution of a Global Menace," *J Infect Dis*, vol. 215, S28-S36, 2017.
25. C. MacDougall, "Penicillins, Cephalosporins, and Other Beta-Lactam Antibiotics," in *Goodman and Gilman's Pharmacologic Basis of Therapies*, 13th ed., McGraw-Hill Education, 2018.
26. M. F. A. Marjani, "Occurrence of ESBL and MBL Acinetobacter Baumannii," *International Journal of Current Research*, 2013.
27. S. O. Meroueh et al., "Ab Initio QM/MM Study of Class A Beta-Lactamase Acylation: Dual Participation of Glu166 and Lys73 in a Concerted Base Promotion of Ser70," *J Am Chem Soc*, vol. 127, 2005.
28. E. L. Miller, "The Penicillins: A Review and Update," *J Midwifery Women's Health*, vol. 47, pp. 426-434, 2002.
29. C. S. Miller et al., "Short-Read Assembly of Full-Length 16S Amplicons Reveals Bacterial Diversity in Subsurface Sediments," 2013.
30. J. M. Munita and C. A. Arias, "Mechanisms of Antibiotic Resistance," *Microbiol. Spectrum*, 2016.
31. S. D. Oliveira et al., "Mechanisms of Antibacterial Resistance: Shedding Some Light on These Obscure Processes," in *Antibiotic Resistance Mechanisms and New Antimicrobial Approaches*, Academic Press, an

- imprint of Elsevier, 2016.
32. I. Palzkill, "Structural and Mechanistic Basis for Extended-Spectrum Drug-Resistance Mutations in Altering the Specificity of TEM, CTX-M, and KPC Beta-Lactamases," 2018.
 33. K. M. Papp-Wallace et al., "Carbapenems: Past, Present, and Future," 2011.
 34. F. J. Perez-Llarena and G. Bou, "Beta-Lactamase Inhibitors: The Story So Far," 2009.
 35. W. A. Petri, "Penicillins, Cephalosporins, and Other Beta-Lactam Antibiotics," in Goodman and Gilman's the Pharmacological Basis of Therapeutics, 11th ed., USA: McGraw Hill, pp. 1127-1152, 2006.
 36. P. Pottinger et al., "Antibacterial Agents and Resistance," in Sherris Medical Microbiology, McGraw-Hill Education, pp. 435, 437, 444, 2018.
 37. S. Rahman et al., "The Growing Genetic and Functional Diversity of Extended Spectrum Beta-Lactamases," Biomed. Res. Int., 9519718, 2018.
 38. O. Skold, "Antibiotics and Antibiotics Resistance," John Wiley & Sons, Inc., pp. 83, 88, 2011.
 39. T. Strateva and D. Yordanov, "Pseudomonas Aeruginosa - a Phenomenon of Bacterial Resistance," J Med Microbiol, vol. 58, pp. 1133-1148, 2009.
 40. K. S. Thomson, "Extended-Spectrum-Beta-Lactamase, AmpC, and Carbapenemase Issues," J Clin Microbiol, vol. 48, pp. 1019-1025, 2010. <https://doi.org/10.1128/JCM.00219-10>
 41. C. Tooke et al., "Beta-Lactamases and Beta-Lactamase Inhibitors in the 21st Century," Journal of Molecular Biology, 2019. <https://doi.org/10.1016/j.jmb.2019.04.002>
 42. G. Wang et al., "CTX-M Beta-Lactamase-Producing Klebsiella Pneumonia in Suburban New York, New York, USA," Emerging Infectious Diseases, vol. 19, no. 11, pp. 1803, 2013. DOI: <http://dx.doi.org/10.3201.1911.121470>
 43. R. Wax et al., "Bacterial Resistance to Antimicrobials," Taylor & Francis Group, LLC, 2nd ed., pp. 122, 114, 2008.
 44. A. F. Wendel et al., "Genetic Characterization and Emergence of the Metallo-Beta-Lactamase GIM-1 in Pseudomonas spp. and Enterobacteriaceae During a Long-Term Outbreak," Antimicrob. Agents Chemother., vol. 57, pp. 5162-5165, 2013
 45. U. U. Zango, M. Ibrahim, and S. A. A. Shawai, "A Review on Beta-Lactam Antibiotic Drug Resistance," MOJ Drug Des Develop Ther., vol. 3, no. 2, pp. 52-58, 2019. DOI: 10.15406/mojddt.2019.03.00080
 46. G. G. Zhanel, R. Love, and H. Adam, "Tedizolid: A Novel Oxazolidinone with Potent Activity Against Multidrug-Resistant Gram-Positive Pathogens," 2015.