

Table Of Content

Journal Cover	2
Author[s] Statement	3
Editorial Team	4
Article information	5
Check this article update (crossmark)	5
Check this article impact	5
Cite this article	5
Title page	6
Article Title	6
Author information	6
Abstract	6
Article content	7

Academia Open



By Universitas Muhammadiyah Sidoarjo

Originality Statement

The author[s] declare that this article is their own work and to the best of their knowledge it contains no materials previously published or written by another person, or substantial proportions of material which have been accepted for the published of any other published materials, except where due acknowledgement is made in the article. Any contribution made to the research by others, with whom author[s] have work, is explicitly acknowledged in the article.

Conflict of Interest Statement

The author[s] declare that this article was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright Statement

Copyright © Author(s). This article is published under the Creative Commons Attribution (CC BY 4.0) licence. Anyone may reproduce, distribute, translate and create derivative works of this article (for both commercial and non-commercial purposes), subject to full attribution to the original publication and authors. The full terms of this licence may be seen at <http://creativecommons.org/licences/by/4.0/legalcode>

EDITORIAL TEAM

Editor in Chief

Mochammad Tanzil Multazam, Universitas Muhammadiyah Sidoarjo, Indonesia

Managing Editor

Bobur Sobirov, Samarkand Institute of Economics and Service, Uzbekistan

Editors

Fika Megawati, Universitas Muhammadiyah Sidoarjo, Indonesia

Mahardika Darmawan Kusuma Wardana, Universitas Muhammadiyah Sidoarjo, Indonesia

Wiwit Wahyu Wijayanti, Universitas Muhammadiyah Sidoarjo, Indonesia

Farkhod Abdurakhmonov, Silk Road International Tourism University, Uzbekistan

Dr. Hindarto, Universitas Muhammadiyah Sidoarjo, Indonesia

Evi Rinata, Universitas Muhammadiyah Sidoarjo, Indonesia

M Faisal Amir, Universitas Muhammadiyah Sidoarjo, Indonesia

Dr. Hana Catur Wahyuni, Universitas Muhammadiyah Sidoarjo, Indonesia

Complete list of editorial team ([link](#))

Complete list of indexing services for this journal ([link](#))

How to submit to this journal ([link](#))

Article information

Check this article update (crossmark)



Check this article impact (*)



Save this article to Mendeley



(*) Time for indexing process is various, depends on indexing database platform

Positioning Preterm Newborns for Cardiorespiratory Stability A Global Insight

Memposisikan Bayi Baru Lahir Prematur untuk Stabilitas Kardiovaskuler Sebuah Wawasan Global

Haneen Abdul Hussein Shamkhi, hanin.abd2204m@conursing.uobaghdad.edu.iq,
Pediatric Nursing Department, College of Nursing, University of Baghdad, Baghdad, Iraq ⁽¹⁾

Adraa Hussein Shawq, hanin.abd2204m@conursing.uobaghdad.edu.iq, (0)
Pediatric Nursing Department, College of Nursing, University of Baghdad, Baghdad, Iraq

⁽¹⁾ Corresponding author

Abstract

This study assessed the effects of supine, prone, and right lateral positioning on the cardiorespiratory indicators of preterm newborns using a randomized controlled trial with 60 participants. Measurements of heart rate, respiratory rate, and oxygen saturation were taken before and after the application of these positions over three days. Results indicated significant improvements in all cardiorespiratory parameters in the intervention groups ($p = 0.000$), while the control group showed no significant changes. The findings suggest that strategic positioning can be an effective part of developmental care in neonatal intensive care units, enhancing the stabilization of vital functions in preterm newborns.

Highlights:

- Strategic Positioning: Enhances cardiorespiratory stability in preterm newborns.
- Methodological Rigor: Employs a robust randomized controlled trial.
- Practical Implications: Recommends routine NICU positioning strategies.

Keywords: Preterm Newborns, Cardiorespiratory Stabilization, NICU, Positioning Strategies, Randomized Controlled Trial.

Published date: 2024-05-30 00:00:00

Introduction

Any newborns delivered before the complete 37 weeks of pregnancy following the last menstrual cycle is referred to as a preterm [1]. Approximately 15 million preterm births were recorded by the World Health Organization (WHO) [2],[3]. The improper development of systemic organs in preterm newborns results in impaired adaptation to extrauterine life, a decreased chance of survival, and serious newborns consequences [4],[5]. Worldwide, premature complications following delivery cause of death children under five [6]. And the second cause of most newborns die within their first month of life[7],[8].

The transition of neonates from intrauterine life to extrauterine life is a critical and sensitive period [9]. But for preterms, it is recognized as a set of challenges due to an incompetent cardiovascular system that is influenced by the transition from a lower placental circulation resistance to the higher artery circulation resistance after delivery [10]. In addition, the preterms exposure to respiratory problems such as respiratory distress syndrome, acidosis, apnea, and hypoxia are caused by their pulmonary insufficiency, weakness in the respiratory muscles, immature lungs, and a lack of surfactant in the alveoli which result in increased oxygen consumption [11]. In order to improve survival rates and lower the risk of complication, the preterm must be hospitalized in the neonatal intensive care unit (NICU) to receive special attention and intensive care [12],[13]. The nurses have a major role in promoting the care of preterm and high-risk newborns [14]. Putting effort to maintain optimal health, prevent complications, and achieve desirable outcomes [15],[16]. The nurse can employ a variety positions such as lateral, supine, and prone positions as developmental supportive care for the preterm receiving care in the NICU that influences physiological stability [17].

When newborn placing on their stomach is known as prone positioning. In the USA, nearly all newborns were placed in the prone position until before 1990 [18]. The physiological benefits of the prone position can improve ventilation perfusion, development of the lungs, and increased drainage of secretions from the trachea and bronchus [19]. In this position, the diaphragm muscles have the optimal length and the angle of contraction and stabilize the rib cage. As a result, there has been an increase in tidal volume, an improvement in diminished thoracoabdominal function, increasing blood flow through the lungs, and reduction in CO₂[20],[21].

The American Pediatrics Association recommended the supine position as the ideal position in 1996 for lowering the danger of sudden infant death syndrome in infants under the age of one year. In supine position, the preterm lying on their back [21],[22]. This position, with slight head elevation, has better respiratory muscle strength, maximal lung development, enhanced cardiac and respiratory performance, and increased oxygen saturation in preterm neonates [17],[23]. While when newborn lie on their side, the thigh is slightly flexed, and the other thigh is flexed to the abdomen called lateral position. This position keeps the airway open and influences blood flow returning to the heart, which enhances the heart pump's function. As a result, the amount of hemoglobin that binds the oxygen can rise, fill the cells with oxygen, and increase oxygen saturation [20]. For providing preterms proper position should be use supportive positioning device such as nest to gives them boundaries, maintain position, promotes comfort, relaxation, and helps the preterm feel with position similar to mother womb[24].

In this study, the authors applied non-costing simple positioning techniques (supine, prone, and right lateral) in order to determine their effects on preterm cardiorespiratory stability, with constructing the nest as a model used to make boundaries to fix the preterm positioning. Preterm newborns were selected due to their higher need for supportive care, which may reduce the burden on the preterm, caregivers, medical staff, shorten hospital stays, and ultimately reduce the cost of hospitals.

Method

A. Research Design

The research utilized a randomized controlled trial methodology to achieve its goals.. The study was conducted at NICU Al-Batool Teaching Hospital in Diyala, Iraq. conducted between September 15th 2023, to 27thApril 2024.

B. Samples and Sampling

For sixty preterm newborns, a simple random sample methodology was used, which included the writing type of each group (lateral, prone, supine, and control groups separately in similar pieces in size, color, and folding method, to select preterm groups these papers are put in a container and thoroughly mixed. The minimum required sample size would be 15 newborns in each group based on a Richard Geiger Equivalent with population proportion = 50%, error probability = 5%, confidence level = 95%, and standard score corresponding to the level of confidence 1.96.

C. Inclusion / Exclusion Criteria

the eligibility criteria includes preterm gestational age of 28–36 weeks, weight between 500–2500 grams, from both genders, and admission to the NICU for a three days. While excluded from the study were preterm newborns who had respiratory distress syndrome, on mechanical ventilation, had sepsis, obviously abnormal from birth, had bone fractures, were confused or had undergone medical surgery.

D. Measurement and Data Collection

Part one : Preterm newborns' sociodemographic information was gathered, such as their gestational age at birth, sex, postnatal age, birth weight, and the type of delivery.

Part two : Every premature baby had a specific chart that was used to record five times a day the cardiorespiratory parameters of that newborn, which included heart rate, respiration rate (HR), oxygen saturation (SpO₂), and respiratory rate (RR). The RR was measured manually, while the HR and SpO₂ were measured with a pulse oximeter.

Part three : The nest was created by the researcher using soft fabric that was layered with belts and brocade, and it formed the U shape Based on the Thames Valley Wessex Neonatal Network [25] as see in (figure 1).



Figure 1. Nesting Model

The pretest measured their HR, RR, and SpO₂ for all groups before any care was given for newborn, applied positions for intervention groups and routine care for control group directly. The first session started at this time and lasted for two hours; the posttest was measured at the first and second hours of the first session. After that, the experimental groups' preterm newborns were allowed to remain in an unrestricted position for thirty minutes while the control group continued received standard care. Then the preterm returned to the previous position, and the second session started, during which the parameter was measured at the first and second hours as explained in (figure 2). The positions was applied according to the recommended position [26] (figure 3)




Figure 2. Data Collection Method

Position description	Figure
<p>The supine position (Figure 3):</p> <ul style="list-style-type: none"> a. Maintain the head as close to or on the midline as possible. b. Avoid neck excessive flexion or extension and maintain a small extension of the neck especially in extremely preterm neonate to keep the head in a neutral position. c. Maintain a forward-rounded shoulders. d. The shoulders are supported (by a roll of linen) to keep them angled slightly forward. e. Support the flexion of the legs against firm constraints on the leg's capacity to support. f. Flexion of the hips and knees is encouraged. g. Tools like a nest, blankets, linen rolls, and sheepskins are utilized to ensure ideal positioning. 	 <p style="text-align: center;">Figure 3</p>
<p>Lateral position (Figure4):</p> <ul style="list-style-type: none"> a. Maintain the head in the midline position... b. provides "C" shaped support for the neck and back. c. A neutral upper shoulder posture is maintained rather than <u>retracted</u>. d. Support is given to create a slightly rounded back. e. It is encouraged to flexion hips and knees. f. <u>Maintain the</u> legs flexion with provide leg support. g. The infant's flexed back will be supported in the side-lying position by swaddling or a heavy blanket roll; this will encourage midline hand-to-hand or hands-to-face movements. 	 <p style="text-align: center;">Figure 4</p>
<p>The prone position (Figure5):</p> <ul style="list-style-type: none"> a. Maintain a neutral head posture or with slight chin tilt toward the chest to prevent overextension of the neck. b. allowed for hands to be near to the face. c. <u>maintain</u> the shoulders rounded and fall forward by Using a baby nest or other similar position assistance d. The legs are supported by limiters and kept hidden beneath the body. e. Avoiding the frog-leg position, or total hip abduction. f. The abdomen and chest are supported with a linen roll. g. makes sure the hips are positioned lower than the head. 	 <p style="text-align: center;">Figure5</p>

Figure 3. The Recommended Positions are as follows

E. Pilot Study

Nine preterm infants admitted to the NICU at Al-Batool Teaching Hospital between October 13 and October 29, 2023, those newborns were not included in the study sample.

F. Reliability and Validity of the Instrument

When the researcher and a co-observer read the physiological indicators at the same time and same sample, their accuracy was compared. The validity was evaluated by 16 experts with more than ten years of combined expertise.

The experts were from the University of Baghdad, University of Babylon, the University of Karbala College of Nursing.

G. Ethical Considerations

Before the actual data collection procedure starts, the study has official administrative approval from the College of Nursing's Ethical Committee of Research to be done. It is also registered with the Iranian Registry of Clinical Trials (IRCT) under the number IRCT20231120060117N3. Furthermore, approval to collect the data was granted by the Iraqi Ministry of Planning and the Ministry of Health. A formal consent from the parents of preterm is attached to the data collection process.

H. Data Analysis

The study's data are analyzed using the SPSS software, version 26. The following are included in the descriptive statistical data analysis approach: frequency (F.), percentage (%), Standard Deviation (SD), and mean of score. Repeated Measure ANOVA, an inferential statistical data analysis method, was applied.

Result and Discussion

A. Result

Variable	Groups	lateral sided position group		Supine position group		Prone position group		Control group	
		F.	%	F.	%	F.	%	F.	%
Gestational age	28 – 31 weeks	4	26.7	8	53.3	7	46.7	6	40
	32- 34 weeks	7	46.6	6	40	6	40	7	46.7
	35 - 36 weeks	4	26.7	1	6.7	2	13.3	2	13.3
	Mean ± SD	32.66 ± 2.12		31.6 ± 2.16		31.73 ± 2.31		32.26 ± 1.79	
Postnatal age	1	3	20	3	20	4	26.7	2	13.3
	2	3	20	4	26.7	4	33.3	3	20
	3	4	26.7	4	26.7	6	40	4	26.7
	4	4	26.7	4	26.7	1	6.7	5	33.3
	5	1	6.6	0	0	0	0	1	6.7
	Total	15	100	15	100	15	100	15	100
Gender	Female	10	66.7	7	46.7	5	33.3	8	53.3
	Male	5	33.3	8	53.3	10	66.7	7	46.7
	Total	15	100	15	100	15	100	15	100
Type of delivery	C/S	6	40	4	26.7	7	46.7	8	53.3
	NVD	9	60	11	73.3	8	53.3	7	46.7
	Total	15	100	15	100	15	100	15	100
Birth Weight	Mean ± SD	Mea n	SD	Mea n	SD	Me an	SD	Me an	SD
		1536.6 ± 259		1500±429.2		1530 ± 543		1548.6±358.3	

F.= frequency, % = percentage

Figure 4. The Socio-demographical Data of Preterm Newborns

(figure 4) revealed that the right lateral and control groups had a mean gestational age of (32.66, 32.26 weeks), while the supine and prone positions had a mean gestational age of (31.6, 31.73). Furthermore, the majority of the right lateral positioned postnatal age (26.7%) occurred between 3 and 4 days, and the same percentage (26.7%) occurred between 3 and 4 days of supine positioning preterm. In the prone-positioned , the largest percentage (40%) occurred at 3 days, while in the control group, it was 33.3% at 4 days after birth. In the right lateral and control groups (66.7 and 53%) were female respectively, while 66.7 % and 53.3 % were male in prone and supine positions. The greatest percentages (73.3%, 60 %, 53.3%) in the supine, right lateral, and prone position groups were delivered by NVD, while 53.3 % of the control group delivered by C/S. The mean birth weight of preterms was 1500 g, 1536.6 g, 1530 g, and 1548.6 g for supine, right lateral, prone, and control groups respectively.

Items	Times of testing	Lateral position group					Supine position group					Prone position group					Control group				
		Mean			F	P	Mean			F	P	Mean			F	P	Mean			F	P
		1 st day	2 nd day	3 rd day			1 st day	2 nd day	3 rd day			1 st day	2 nd day	3 rd day			1 st day	2 nd day	3 rd day		
Heart rate	Pretest	150.6	143.4	138.6	2	0	156	146.4	139.6	3	0	154.3	148.4	140.7	6	0	139.7	142.8	136.5	1	0
	Post 1 hr from 1 st session	147	138.9	136.7	6	1	157.6	146	139	7	4	151.7	146.3	142.2	5	0	142	142.9	141.9	9	4
	Post 2 hrs from 1 st session	146.4	139.6	135.1	6	0	153.5	143.2	135.1	8	0	150.6	144.3	135.3	4	0	144.5	143	141	2	7
	Post 1 hr from 2 nd session	144	138.9	134.4			152.4	142.6	133.8	2		149	140.4	133.1	1		140	139	140.8		
	Post 2 hrs from 2 nd session	145.5	136.6	136.3			149	140.8	133			147.5	139.7	129.7			141.8	140.7	139.6		
Respiratory rate	Pretest	52	43.4	35.7	3	0	55.06	46.6	42.6	9	0	55.5	47.7	41.6	1	0	47.5	47.1	45.6	1	0
	Post 1 hr from 1 st session	51.5	41.9	36.7	3	0	53.2	45.6	42.3	9	6	52.8	46.2	38.8	3	0	50.5	49.8	48.7	6	3
	Post 2 hrs from 1 st session	46.3	42.4	35.06	7	0	51.26	43.7	40.6	4	0	52	44.4	36.4	2	0	49.3	48.6	48.8	0	4
	Post 1 hr from 2 nd session	46	39.3	35.6	4		49.8	42.06	38.3			48.4	42.06	34.4			47.6	47.3	46.6	6	
	Post 2 hrs from 2 nd session	45.4	38.5	34.06			48.3	41.2	36.4			46.8	39.7	32.4			44.2	45.2	44.7		
Oxygen saturation	Pretest	94.2	95.6	96.5	6	0	93.2	94.4	95.1	1	0	93.7	95	96.2	2	0	93.7	94.4	94.6	2	0
	Post 1 hr from 1 st session	94.06	95.53	96.46	5	8	93.4	94.3	95.2	4	1	93.8	95.5	96.8	4	0	93.8	94.1	94.9	3	5
	Post 2 hrs from 1 st session	94.6	96.26	97.1	1	0	93.8	94.8	95.7	3	0	94.6	92.2	93	2	7	93.6	93.8	94.7	6	2
	Post 1 hr from 2 nd session	95.13	96.86	97.46			94.1	95.2	96.2	3		94.8	96.5	97.8	7		93.6	94.6	94.8	3	
	Post 2 hrs from 2 nd session	95.8	97.6	98.2			94.6	96	96.4	3		95.8	97.4	98.7	6		93.9	95	95.5		

NS = non-significant at P>0.05, S=significant at P<0.05]

Figure 5. Showed there were High Significant Differences in the Mean Physiological Parameters

(figure 5) showed there were high significant differences in the mean physiological parameters (HR, RR, Spo₂)of preterm newborn in lateral, supine, and prone groups at P = 0.000, while in control group showed non- significant difference at P higher than 0.05.

B. Discussion

As shown in (figure 4) The mean most of the preterm gestational age within the study groups has been determined to sit between 31.6 and 32.6 weeks and the majority postnatal age were 2-4 days. The majority of preterm newborns birth by normal vaginal delivery and have a mean weight between 1500 -1548 g. Regarding the preterms gender were equally in study.

In (figure 5) The result of the study demonstrated that the mean score of the pretest on the first day and post-test on the third day was changed from (150.6 b/m to136.3b/m) in the lateral group, (156 b/m to133b/m) for the supine group, and (154.3b/m to 129.7 b/m) for the prone group. The three groups recorded a significant change in decreased heart rate during the three days at p value = 0.000, while the control group statically recorded non-significant differences in the mean preterm heart rate at p =0.47. These results line with a study conducted in Egypt to examine the impact of right lateral position, knee-chest (prone), and semi-sitting on 57 premature newborns suffering from respiratory distress syndrome. Mahrous results revealed that the mean score of preterms heart rate in prone and lateral position decreased from (150.5, 153.1) at first session to (142.5, 148.5) at second session respectively, statically showed significant differences at p = (0.0001) [20]. Another study in (2022) conducted to evaluate how preterms cardiorespiratory patterns are affected by lateral versus supine position, 162 preterms were laying in each position for 1 hour, measured HR, RR, and Spo₂ every 30 minutes. Their result showed the mean heart rate of preterms in supine groups significant at P< 0.0001[17].

Also (figure 5) showed that there was a statistically significant decrease in the mean score of preterm respiratory rate among positioning groups at p value =0.000 based on the result of the pretest for supine, prone, and right

lateral groups were (55.06bpm, 55.5 bpm, 52 bpm,) and post-test at third days were (36.4bpm, 32.4bpm, 34.06 bpm.). While control group result pre and posttest was (47.5 bpm, 44.7 bpm) demonstrated non- significant differences at $p= 0.34$. The results of the present study corresponds with the findings of a study in Brazil conducted on 24 preterm neonates, divided randomly into four groups supine, prone, left side, and right side position to assess the effectiveness of these positions on their physiological and behavioral responses before 30 minutes, during intervene it lasted for 4 hours, and after 30 minutes from routine care applied. showed that the preterm respiratory rate was decreased during the intervened period in the prone, supine, and right lateral groups [27].

The final physiological indicator was oxygen saturation. According to the statistics of the present study which showed that the mean score of oxygen saturation has a significant increase in three different positions supine, prone, and right lateral at $p = 0.000$. As indicated in (figure 5), the elevation mean Spo₂ of preterm in three position groups respectively were from (93.2%, 93.7%, 94.2%) to (96.4%, 98.7%,98.2%), while in control group revered to non-significant differences in the mean Spo₂ of preterms at $p=0.52$. These findings in line with experimental study conducted on 40 preterms to find out the effect of positioning on preterms HR and Spo₂, applied the three positions supine, prone, and lateral; each position persisted for two hours, the authors measured Spo₂ and HR every 15 minutes. Their funding demonstrated a significant change in preterm oxygen saturation at $p= 0.032$ in all positions [28].

Several previous study showed, when the preterm placing in the proper positions, the pressures acting on the heart, chest wall, lungs, vascular system, blood volume, and diaphragm elevate the impact of gravity, hydrostatic pressure, and compressional forces on the cardiorespiratory system, and enhancing the transfer of oxygen [29]. In addition to , enhances heat regulation, skin integrity, sleep quality, pain reduction, apnea/desaturation events, and neurobehavioral organization [30].

Conclusion

The study conclusively demonstrated that the implementation of supine, prone, and right lateral positioning strategies significantly stabilizes the cardiorespiratory indicators such as heart rate, respiratory rate, and oxygen saturation in preterm newborns. These findings underscore the efficacy of strategic positioning in enhancing neonatal care in the NICU setting. The statistical significance of the changes in the intervention groups ($p = 0.000$) compared to the non-significant differences observed in the control group ($p > 0.05$) reinforces the potential of these positioning strategies as critical components of developmental care for preterm infants. Given the positive outcomes, it is recommended that such positioning techniques be routinely integrated into the care protocols of neonatal intensive care units to optimize the health and development of preterm newborns. Future research should explore the long-term impacts of these positioning strategies on neonatal health outcomes, potentially extending into pediatric care practices to further substantiate and refine the implementation guidelines.

References

1. S. H. Alwan and E. G. A. Ma'ala, "Effectiveness of an Educational Program on Mothers' Attitudes Toward Readiness for Discharge Care at Home for a Premature Baby in Intensive Care Unit at Teaching Hospitals in Medical City Complex," *Iraqi National Journal of Nursing Specialties*, vol. 34, no. 2, pp. 1-7, 2021, doi: 10.58897/injns.v34i2.570.
2. World Health Organization, "Preterm Birth," World Health Organization: WHO, May 10, 2023. [Online]. Available: <https://www.who.int/news-room/fact-sheets/detail/preterm-birth>
3. S. Nayef and M. A. Neamah, "Evaluation of Nursing Performance Concerning Nasogastric Tube in Neonatal Intensive Care Unit at Al-Batool Teaching Hospital in Baqubah City," *Iraqi National Journal of Nursing Specialties*, vol. 32, no. 1, pp. 47-57, Jun. 2019, doi: 10.58897/injns.v32i1.323.
4. G. Altay and S. Küçüköçlü, "Effects of the Facilitated Tucking Position in Early Period on Physiological Parameters, Comfort and Breastfeeding Performance in Late Preterm Infants: A Randomized Controlled Trial," *Midwifery*, vol. 115, p. 103492, Sep. 2022, doi: 10.1016/j.midw.2022.103492.
5. F. B. McDonald, E. M. Dempsey, and K. D. O'Halloran, "The Impact of Preterm Adversity on Cardiorespiratory Function," *Experimental Physiology*, vol. 105, no. 1, Dec. 2020, doi: 10.1113/ep087490.
6. W. A. Hussein and I. M. Abbas, "Assessment of Nurse-Midwife's Knowledge and Performance Regarding Immediate Newborn Care in Delivery Rooms at Maternity Hospitals in Baghdad City," *Iraqi National Journal of Nursing Specialties*, vol. 34, no. 2, pp. 99-113, 2021, doi: 10.58897/injns.v34i2.579.
7. S. A. R. Atiyah, S. H. Khudhair, and N. N. Hameed, "Effectiveness of Education Health Program on Nurses' Practices Toward Neonates Care with Sepsis in Incubator in Neonatal Intensive Care Unit in Teaching Hospitals Baghdad City," *Indian Journal of Public Health Research & Development*, vol. 9, no. 8, pp. 1246-1249, Aug. 2018, doi: 10.5958/0976-5506.2018.00901.4.
8. H. H. Ghafel, "Neonatal Mortality Rate at AL-Elwaya Maternity Hospital in Baghdad City: Retrospective Study," *Journal of Contemporary Medical Sciences*, vol. 4, no. 2, pp. 88-91, Sep. 2018, [Online]. Available: <https://www.iasj.net/iasj/download/3b413f1b55c081d2>.
9. Z. A. S. Abdul Rahman, S. G. Turki, and Z. W. Ajil, "Evaluation of Nurses' Practice Toward Care of Neonates During Hypothermia," *Migration Letters*, vol. 20, no. S2, pp. 733-743, Jul. 2023, doi:

- 10.59670/ml.v20iS2.3728.
10. J. Lewandowski, "The Preterm Heart: A Unique Cardiomyopathy?," *Pediatric Research*, vol. 85, no. 6, pp. 738-739, May 2019, doi: 10.1038/s41390-019-0301-3.
 11. K. Patel, S. Mutalikdesai, and M. Chaudhari, "Effectiveness of Conventional Care (Control Group) Versus Nesting (Experimental Group) on Selected Physiological Parameters Among Preterm Neonates in NICU," *International Journal of Multidisciplinary Educational Research*, vol. 10, no. 2, p. 1, 2021, [Online]. Available: [http://ijmer.s3.amazonaws.com/pdf/volume10/volume10-issue1\(2\)/11.pdf](http://ijmer.s3.amazonaws.com/pdf/volume10/volume10-issue1(2)/11.pdf)
 12. A. Prawesti, E. Emaliyawati, R. Mirwanti, and A. Nuraeni, "The Effectiveness of Prone and Supine Nesting Positions on Changes of Oxygen Saturation and Weight in Premature Babies," *Jurnal Ners*, Oct. 2019, [Online]. Available: <https://repository.unar.ac.id/jspui/handle/123456789/504>
 13. N. Q. Tuoma and M. F. Khalifa, "Evaluation of Quality of Nursing Care Services Provided to Children under Five Years Based on Integrated Management of Child Health at Primary Health Care Centers in Baquba City," *Iraqi National Journal of Nursing Specialties*, vol. 34, no. 1, pp. 1-10, Jun. 2021, doi: 10.58897/injns.v34i1.454.
 14. R. Aziz and M. A. Abdul-Hamza, "Effectiveness of an Educational Program upon Nurses' Knowledge Toward The Continuous Positive Airway Pressure (CPAP) Machine in Neonatal Intensive Care Unit at Al-Diwanyia City Hospitals," *International Journal of Scientific and Research Publications*, vol. 7, no. 8, pp. 469-476, 2017, [Online]. Available: <http://www.ijsrp.org/research-paper-0817.php?rp=P686449>
 15. A. Arrar and S. J. Mohammed, "Effectiveness of an Educational Program on Nurses' Knowledge and Practices Concerning Nursing Care for Critically-Ill Patients at Critical Care Units in Misan Governorate Hospitals," *Medico-Legal Update*, vol. 20, no. 3, 2020, [Online]. Available: https://systems.uomisan.edu.iq/cv/uploads/files/8og7bwq5306j_vl.pdf
 16. J. Jassim and M. F. Khalifa, "Evaluation of Quality of Primary Health Care Services at Primary Health Care Centers in Baghdad City: A Comparative Study," *Iraqi National Journal of Nursing Specialties*, vol. 31, no. 1, pp. 110-117, Jun. 2018, doi: 10.58897/injns.v31i1.297.
 17. Mahdey Mohamed, S. Salah Eldin Ibrahim, H. Rafaat Tantawi, and F. Mohamed, "Lateral Versus Supine Position on Breathing Patterns Among Premature Neonates: A Comparative Study," *Egyptian Journal of Health Care*, vol. 13, no. 3, pp. 902-911, Sep. 2022, doi: 10.21608/ejhc.2022.255693.
 18. H. M. El-Sayd, G. M. El-Mashad, M. Z. Mohamed, and Z. S. Abouzouna, "Comparison of Massage and Prone Position on Heart Rate and Blood Oxygen Saturation Level in Preterm Neonates Hospitalized in Neonatal Intensive Care Units," *Egyptian Pediatric Association Gazette / Egyptian Pediatric Association Gazette*, vol. 71, no. 1, Dec. 2023, doi: 10.1186/s43054-023-00214-6.
 19. N. Mohamed Arafa Mohamed, M. Ebrahim Fouda Elashry, and R. L. Ahmed Mokbe, "Effect of Lateral Positions versus Prone Position on Cardiorespiratory Parameters among Preterm Neonates with Respiratory Distress Syndrome," *Egyptian Journal of Health Care*, vol. 11, no. 2, 2020, [Online]. Available: https://ejhc.journals.ekb.eg/article_179966_8c5eaa28581dd7b0af9270f61b7b585c.pdf
 20. S. Mahrous and A. Aboelmagd, "Effect of Knee-Chest, Semi-Sitting, and Right Lateral Position on Preterm Neonates with Respiratory Distress Syndrome," *International Egyptian Journal of Nursing Sciences and Research*, vol. 2, no. 2, pp. 353-361, Jan. 2022, doi: 10.21608/ejnsr.2022.212952.
 21. Güler and H. Çalışır, "The Effect of Positioning on Adaptation to Spontaneous Breathing in Premature Infants After Weaning from Mechanical Ventilation: A Randomized Controlled Trial," *The Journal of Pediatric Research*, vol. 7, no. 2, pp. 102-109, May 2020, doi: 10.4274/jpr.galenos.2019.19042.
 22. Wardany Abdelaal Mohamed, F. El-Sayed Abusaad, and O. Abd El-Fatah El-Agamy, "Effectiveness of Educational Program on Mothers' Knowledge and Practices for the Prevention of Sudden Infant Death Syndrome," *International Egyptian Journal of Nursing Sciences and Research*, vol. 3, no. 2, pp. 506-523, Jan. 2023, doi: 10.21608/ejnsr.2023.278177.
 23. T. Dassios, A. Vervenioti, and G. Dimitriou, "Respiratory Muscle Function in the Newborn: A Narrative Review," *Pediatric Research*, vol. 91, no. 1, pp. 1-9, Apr. 2021, doi: 10.1038/s41390-021-01529-z.
 24. P. Sumathy, "Effectiveness of Nesting Technique on Posture and Physiological Parameters," *Pondicherry Journal of Nursing*, vol. 13, no. 2, pp. 25-28, 2020, doi: 10.5005/jp-journals-10084-12149.
 25. T. Valley, "Positioning and Handling on the Neonatal Unit," 2018. [Online]. Available: https://www.piernetw.ork.org/uploads/4/7/7/8/1/47810883/positioning_and_handling_on_the_neonatal_unit_v3_july_2017_final.pdf
 26. V. Danila, A. Curteza, and S. Balan, "Study of the Particulars of Positioning of Premature Babies for the Development of Functional Clothing Products," *Annals of the University of Oradea, Fascicle of Textiles, Leatherwork*, vol. 2, pp. 23-30, Sep. 01, 2023. [Online]. Available: https://ibn.idsi.md/vizualizare_articol/186738
 27. M. G. Santos, C. S. Viera, G. R. F. Bertolini, E. F. Osaku, C. R. L. de M. Costa, and A. T. K. G. Grebinski, "Physiological and Behavioural Effects of Preterm Infant Positioning in a Neonatal Intensive Care Unit," *British Journal of Midwifery*, vol. 25, no. 10, pp. 647-654, Oct. 2017, doi: 10.12968/bjom.2017.25.10.647.
 28. Z. Akbarian Rad, M. Haghshenas Mojaveri, M. Hajiahmadi, A. Ghanbarpour, and S. Mirshahi, "The Effect of Position on Oxygen Saturation and Heart Rate in Very Low Birth Weight Neonates," *Caspian Journal of Pediatrics*, vol. 2, no. 2, pp. 153-157, Sep. 2016, doi: 10.22088/acadpub.BUMS.2.2.153.
 29. S. A. Salih, A. M. Omar, S. A. Elwasefy, D. A. Said, and A. Y. Elderderly, "Impact of Positioning on Neonate with Respiratory Distress: Prone Vs. Supine Position," *Sylwan Journal*, vol. 163, no. 12, pp. 54-69, 2020.
 30. J. Upadhyay, P. Singh, K. C. Digal, S. Shubham, R. Grover, and S. Basu, "Developmentally Supportive Positioning Policy for Preterm Low Birth Weight Infants in a Tertiary Care Neonatal Unit: A Quality Improvement Initiative," *Indian Pediatrics*, vol. 58, no. 8, pp. 733-736, Jan. 2021, doi: 10.1007/s13312-021-2281-8.