

Table Of Content

Journal Cover	2
Author[s] Statement	3
Editorial Team	4
Article information	5
Check this article update (crossmark)	5
Check this article impact	5
Cite this article	5
Title page	6
Article Title	6
Author information	6
Abstract	6
Article content	7

Academia Open



By Universitas Muhammadiyah Sidoarjo

Originality Statement

The author[s] declare that this article is their own work and to the best of their knowledge it contains no materials previously published or written by another person, or substantial proportions of material which have been accepted for the published of any other published materials, except where due acknowledgement is made in the article. Any contribution made to the research by others, with whom author[s] have work, is explicitly acknowledged in the article.

Conflict of Interest Statement

The author[s] declare that this article was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright Statement

Copyright © Author(s). This article is published under the Creative Commons Attribution (CC BY 4.0) licence. Anyone may reproduce, distribute, translate and create derivative works of this article (for both commercial and non-commercial purposes), subject to full attribution to the original publication and authors. The full terms of this licence may be seen at <http://creativecommons.org/licences/by/4.0/legalcode>

EDITORIAL TEAM

Editor in Chief

Mochammad Tanzil Multazam, Universitas Muhammadiyah Sidoarjo, Indonesia

Managing Editor

Bobur Sobirov, Samarkand Institute of Economics and Service, Uzbekistan

Editors

Fika Megawati, Universitas Muhammadiyah Sidoarjo, Indonesia

Mahardika Darmawan Kusuma Wardana, Universitas Muhammadiyah Sidoarjo, Indonesia

Wiwit Wahyu Wijayanti, Universitas Muhammadiyah Sidoarjo, Indonesia

Farkhod Abdurakhmonov, Silk Road International Tourism University, Uzbekistan

Dr. Hindarto, Universitas Muhammadiyah Sidoarjo, Indonesia

Evi Rinata, Universitas Muhammadiyah Sidoarjo, Indonesia

M Faisal Amir, Universitas Muhammadiyah Sidoarjo, Indonesia

Dr. Hana Catur Wahyuni, Universitas Muhammadiyah Sidoarjo, Indonesia

Complete list of editorial team ([link](#))

Complete list of indexing services for this journal ([link](#))

How to submit to this journal ([link](#))

Article information

Check this article update (crossmark)



Check this article impact (*)



Save this article to Mendeley



(*) Time for indexing process is various, depends on indexing database platform

Enhanced Protease Production from Diabetic Ulcer-Derived *E. coli*

*Peningkatan Produksi Protease dari *E. coli* yang Berasal dari Ulkus Diabetes*

Mustafa Talib Muhsin, abualshooq44@gmail.com, (1)

College Biotechnology /Molecular and Medical Biotechnology, Iraq

⁽¹⁾ Corresponding author

Abstract

This study explored a local *Escherichia coli* isolate from diabetic foot ulcer patients for protease enzyme production. Qualitative screening on skim milk agar showed a 6 mm hydrolysis halo, while quantitative screening with casein revealed a specific activity of 144.55 U/mg protein. Purification via ion exchange chromatography on DEAE-cellulose increased the specific activity to 767.5 U/mg, with a 5.3-fold enhancement and a 29.5% yield. The findings suggest *E. coli* as a viable source for protease production and demonstrate the efficiency of ion-exchange chromatography for enzyme purification, with potential significant industrial applications.

Highlight:

E. coli sourced from diabetic ulcers shows promise for protease production.
Skim milk agar and casein used for qualitative and quantitative screening.
Ion exchange chromatography significantly enhances enzyme specific activity.

Keyword: *Escherichia coli*, protease production, diabetic foot ulcer, enzyme purification, ion exchange chromatography

Published date: 2024-06-18 00:00:00

Introduction

The biological processes can now successfully rival traditional chemical processing thanks to recent developments in the field of biocatalysts. Utilizing the most desirable characteristics of biocatalysts, or enzymes, such as high selectivity with fewer side effects and increased yields, mild response circumstances, and frequently low environmental consequences, combinations of chemical and biocatalytic systems are being created there. [1] Two thirds of all the enzymes used in different businesses were proteases [2]. It is used in the waste treatment, food, detergent, and pharmaceutical industries, among others [3]. While proteases used in medicine are manufactured in tiny amounts but need to be thoroughly purified, those prepared in large quantities and utilized as raw materials in the food and detergent industries. formulations.established and characterized extracellular proteases showed proteolytic activity when cultured on skim milk plates [4]. identified and characterized extracellular proteases [5].when cultivated on skim milk plates, shown proteolytic activity [6]. Gram-negative E. coli is a facultative anaerobic bacteria that can convert to fermentation or anaerobic respiration in the absence of oxygen. If oxygen When present, aerobic respiration is used to make ATP.The size of cells are typically 2.0 µm in length, 0.25-1.0 µm in width, and 0.6-0.7 µm³ in volume. Typically, cells appear as rods.[7][8]. Escherichia coli, or just E. coli, is a type of bacteria that can be found in food, the environment, and the intestines of both people and animals. The E. Coli bacteria are a diverse group of bacteria. Some forms of E. Coli can cause illness, but most types are not hazardous.[9][10].

Method

Equipment and apparatus

Apparatus	Company/origin Apparatus
Autoclave	Express/Germany
Balance (sensitive)	Ohaus/ Germany
Centrifuge (Portable)	Selectap/Spain
Incubator	Gallenkamp/UK
Laminar air flow	Memmert/Germany
pH meter	Metter-Tolledo/UK
Vortex	Buchi/Switzerland
UV-Visible spectrophotometer	Shemadzu /Japan
Water Distillator	GFL

Figure 1. The instruments and tools utilized in this investigation were as follows: Materials Both Biological and Chemical

Material	Company/origin
Tris- HCl, K ₂ HPO ₄ , KH ₂ PO ₄	BDH/ England
TCA	Sigma/USA
DEAE-Cellulose	Whatman/ England

Figure 2. The research used the following materials.

Media

Medium	Company/Origin
Nutrient agar	Difco/USA
Nutrient broth	Oxoid

Figure 3. Ready-to-use cultural media were all organized following the guidelines provided by industrialized businesses and autoclaved at 121°C, (15 pound/In²) For 900 seconds.

Laboratory prepared media

Skim milk agar medium [11], buffers and solutions, Potassium phosphate buffer solution (pH 7) and It was prepared according to [12]

Protease assay solution

1. 0.1M Tris-HCl buffer with a pH of 8 It was made by dissolving 500 milliliters of distilled water with 6 grams of basic HCl.
2. Solution of casein (1%): One gram of casein was dissolved in one hundred milliliters of 0.1M Tris-HCl buffer solution to create it.
3. 5% Acid trichloroacetic: It was made by dissolving 5g of TCA in an amount of Distilled water ., and then adding Distilled water. to bring the volume up to 100 ml.

Laboratory prepared media

Skim milk is used to make agar medium [13]. Five grams of skim milk were dissolved in fifty milliliters of distilled water, and the mixture was autoclaved for five minutes to create this medium. Next, 50 milliliters of distilled water were used to dissolve 2 grams of agar, and the mixture was autoclaved for 15 minutes. Before being distributed, the medium was well mixed after it had cooled to 45°C. into sterilized Petridis plates, which are then stored at 4°C until needed.

Solutions and buffers

Solution of potassium phosphate buffer, pH 7 [12]. In accordance with [12], it was prepared.

Solution for protease assay

1. 0.1M Tris-HCl buffer with a pH of 8 It was made by dissolving 500 milliliters of distilled water with 6 grams of basic HCl.
2. Solution of casein (1%): To prepare it, 1g of casein was dissolved in 100 ml of 0.1M Buffer solution Tris-HCl.
3. 5% Acid trichloroacetic: This was created by adding D.W. Protein Assay Solution to bring the volume up to 100 ml after dissolving 5g of TCA in a quantity of distilled water.
4. Protein assay solution [14]
5. Bright blue G-250 Coombassie solution.
6. 50 cc of 95% ethanol was used to dissolve 0.1 g of Coombassie Brilliant Blue in order to prepare it.
7. A pH 8.0 Tris-HCl buffer (0.1M).

Methods

Scheme (1) provided a summary of the primary steps in the research strategy, starting with the samples., protease production.




Figure 4.

Scheme(1):mainstep of the research plan

Sterilization Methods [15]

Autoclaving: Moist Heat Sterilization

Autoclaving was used to sterilize the media and solutions for 15 minutes at 121°C (15lb/in2).

1. Sterilization via Heat Using an electric oven, glassware was disinfected for three to five hours at 160 to 180°C.

2. Membrane Filtration for Sterilization Using (0.22) µm millipore filters, heat-sensitive liquids were sterilised by membrane filtration.

Quantitative screening [16]

A fresh culture of bacterial isolate containing 100 µl was added to a conical flask containing nutritious broth, and the mixture was allowed to culture for 24 hours at 30°C. in a shaker incubator spinning at 150 rpm. Following incubation, the pellets were disposed of, the culture was centrifuged, and the supernatants were removed and subjected to a protease activity analysis. To quantify TCA soluble peptide release, protease activity was assessed in triplicate using 1% (w/v) solution of casein. 0.8 milliliters of casein solution and 0.2 milliliters of enzyme solution made up the test combination. It was incubated for thirty minutes at 37°C. . 1 milliliter of TCA was added to halt the reaction, and it was then cooled in an ice bath for ten to fifteen minutes before being centrifuged for twenty minutes at maximum speed. The control was set up with the help of the identical procedures, with the exception that 0.2 milliliters After adding the enzyme solution to the casein solution, the absorbance at 280 nm was determined According to the following equation, enzyme activity is the quantity of enzyme needed in an experimental setting to raise absorbance at 280 nm by 0.01 in a minute.

$$\text{Enzyme activity (U/ml)} = \frac{\text{Absorbance at 280 nm}}{0.01 \times 30 \times 0.2}$$

0.01: Constant
30: Reaction time (min)
0.2: enzyme volume (ml)

$$\text{Specific activity (U/mg)} = \frac{\text{Activity (U/ml)}}{\text{Protein concentration (mg/ml)}}$$

Figure 5. *Protease purification by ion exchange chromatography*

The following procedures for protease purification were followed, as suggested [17][18] According to [21], g of resin were dissolved in 1 liter of distilled water to create a DEAE-Cellulose column. The beads were first given time to settle before being extensively cleaned with D.W. . until they took on a clear look. Using a Buchner funnel, the suspension was filtered through Whattman No. 1 before being discharged. The resin was again suspended in a solution of sodium hydroxide and chloride (0.25 M). The suspension underwent the previously described filtering process once more, followed by multiple washes in a 0.25 M hydrochloric acid solution and D.W, before being adjusted to a pH of 8 using 0.05 M Tris-HCl. Following preparation, the enzyme was added to an equilibrated ion exchange chromatography column. earlier using a pH 8 Tris-HCl buffer at 0.05M. Gradual amounts of sodium chloride (0.1–0.6 M) were used to elute the associated proteins from the column, which was subsequently cleaned using a volume equivalent of the same buffer. Protease was eluted at a flow rate of 30 ml/h (3 ml/fraction). A UV-VIS spectrophotometer set at 280 nm was used to calculate the absorbance of each fraction.

Result and Discussion

Isolation of E.coli E.coli was obtained from Furkan M.Kadhim m.phD student in AI-Nahran University /Bioseparation lab.

Semi- quantitative screening

When the isolate of *E. coli* was grown on agar media with skim milk, forming a hydrolysis halo for the colony allowed for semi-quantitative screens for protease production (Figure 3-1).shown the isolate's ability to hydrolyze skim milk agar media and generate hydrolysis halos. Halos varied in diameter from 6 mm [19]

Quantitative screening

After cultivating in the production broth medium at 37°C for a full day, the local isolate of *E. coli* was quantitatively screened to assess its capacity to produce protease. Protease's particular activity in the crude filtrates was then measured after centrifugation. Findings indicated that the isolate produced proteases. According to [20], In culture filtrates, the specific activity of protease was 144.55 U/mg of protein. . Genetic variations in the genes involved for producing protease account for the variations in the isolates' capacities to produce protease [21].




Figure 6. *E. coli*'s proteolytic activity on 10% skim milk agar following a 24-hour incubation period at 37°C

Purification of protease by Ion exchange chromatography

Negatively charged protease generated by *E. coli* was purified using the ion exchange chromatography method. This method involved applying crude protease to To eliminate all uncharged and positively charged proteins, wash the protease sample in equal volumes with a 0.05 M Tris-HCl buffer solution (pH 8) after equilibrating the column using a DEAE-cellulose column. Next, with gradient sodium chloride concentrations between 0.1 and 0.6 M, the bounded proteins (negatively charged) were eluted.

Figure 2's results demonstrated that whereas three protein peaks emerged following elution by the varying sodium chloride concentrations, only one protein peak was visible throughout the washing step. By calculating each eluted fraction's absorbance at 280 nm, the four protein peaks were found.




Figure 7. A DEAE-Cellulose column (45 x 2 cm) is used in ion exchange chromatography to purify the protease produced by *Escherichia coli*.

Protease activity was measured using the three protein peaks that were discovered during the elution process. The majority of the eluted proteins (fractions 46 to 66) had protease activity, which reached 15.35 U/ml, according to the results. Protein concentration, protease activity, and specific activity were assessed after the fractions that

indicate protease activity were combined and concentrated. Given that protease associated with the anionic ion exchange (DEAE-cellulose), these findings support the notion that protease has a negative net charge. According to table (3-1) results, this concentrate had a protein concentration of 0.02 mg/ml, a specific activity of 767.5 U/mg and a protease activity of 15.35 U/ml. Additionally, it possessed a 29.5% enzyme yield and a 5.3 purification fold [25].

Purification step	Volume (ml)	Enzyme activity (U/ml)	Protein con.(mg/ml)	Specific activity (U/mg)	Total activite(U)	Purification (folds)	Yield(%)
Crude enzyme	100	13.01	0.09	144.55	1300	1	100
DEAE cellulose	25	15.35	0.02	767.5	383.75	5.3	29.5

Figure 8. Purification steps for protease produced by the *E.coli*

Conclusion

1-E.coli can be used as a good source for the fabrication of protease enzyme.

2- The ion-exchange chromatography method of purifying protease on DEAE-cellulose resulted in an enzyme with increased specific activity and purification fold.

References

1. P. Sorup, C. Tils, and O. Wolf, "Biocatalysis: State of the Art in Europe," *The Brookings Papers on Economic Activity*, vol. 3, pp. 783-831, 1998.
2. R. Gupta, C. K. Beg, and P. Lorenz, "Bacterial Alkaline Protease: Molecular Approach and Industrial Application," *Appl. Microbiol. Biotechnol.*, vol. 59, pp. 15-32, 2002.
3. A. Marek and H. K. Sajja, "Strategies for Improving Enzymes for Efficient Biocatalysis," *Food Technol. Biotechnol.*, vol. 42, no. 4, pp. 251-264, 2004.
4. F. Sharmin and M. Rahman, "Isolation and Characterization of Protease Producing Bacillus Strain FS-1," *Agricultural Engineering International*, vol. IX, pp. 1-10, 2007.
5. A. A. Kortt, I. B. Caldwell, J. E. Burns, R. E. Guthrie, and D. J. Stewart, "Characterization and Substrate Specificity of the Extracellular Proteases of *Bacteroides Nodosus* from Virulent and Benign Ovine Isolates," in *Footrot in Ruminants. Proceedings of a Workshop*, Melbourne, 1985, D. J. Stewart, J. E. Petersen, N. M. McKern, and D. L. Emery, Eds. Glebe, NSW: CSIRO Division of Animal Health and Australian Wool Corporation, 1986, pp. 237-243.
6. G. G. Lilley, D. J. Stewart, and A. A. Kortt, "Amino Acid and DNA Sequences of an Extracellular Basic Protease of *Dichelobacter Laodostrs* Show that it is a Member of the Subtilisin Family of Proteases," *Eur. J. Biochem.*, vol. 210, pp. 13-21, 1992.
7. "Facts about *E. coli*," *Dimensions, Bacteria: Diversity of Structure of Bacteria*, Britannica Online Encyclopedia, 2015. [Online]. Available: <http://www.britannica.com>. [Accessed: 25-Jun-2015].
8. A. C. Yu, J. F. Loo, S. Yu, S. K. Kong, and T. F. Chan, "Monitoring Bacterial Growth Using Tunable Resistive Pulse Sensing with a Pore-Based Technique," *Appl. Microbiol. Biotechnol.*, vol. 98, no. 2, pp. 855-862, 2014.
9. O. Tenaillon, D. Skurnik, B. Picard, and E. Denamur, "The Population Genetics of Commensal *Escherichia Coli*," *Nature Rev. Microbiol.*, vol. 8, no. 3, pp. 207-217, 2010.
10. R. L. Vogt and L. Dippold, "Escherichia Coli O157 Outbreak Associated with Consumption of Ground Beef, June-July 2002," *Public Health Reports*, vol. 120, no. 2, pp. 174-178, 2005.
11. P. H. A. Sneath, N. S. Mair, M. E. Sharp, and J. G. Holt, *Bergey's Manual of Systematic Bacteriology*. U.S.A: Williams and Wilkins, 1986, pp. 1104-1138.
12. R. M. Atlas, L. C. Parks, and A. E. Brown, *Laboratory Manual of Experimental Microbiology*. USA: Mosby-Year Book, Inc, 1995.
13. P. H. A. Sneath, N. S. Mair, M. E. Sharp, and J. G. Holt, *Bergey's Manual of Systematic Bacteriology*. U.S.A: Williams and Wilkins, 1986, pp. 1104-1138.
14. M. Bradford, "A Rapid and Sensitive Method for the Quantitation of Microgram Quantities of Protein Using the Principle of Protein-Dye Binding," *Anal. Biochem.*, vol. 72, pp. 248-254, 1976.
15. C. H. Colline and P. M. Lyne, *Microbiological Methods*, 5th ed. London: Butterworth and Co. Publisher Ltd., 1987.
16. P. L. Manachini, M. G. Fortina, and C. Parini, "Production of Proteolytic Enzymes by *Bacillus Acidocaldarius*," *Appl. Microbiol. Biotechnol.*, vol. 28, pp. 409-413, 1989.
17. P. Secades and J. A. Guijarro, "Purification and Characterization of an Extracellular Protease from the Fish Pathogen *Yersinia Ruckeri* and Effect of Culture Conditions on Production," *Appl. Environ. Microbiol.*, vol. 65, pp. 3969-3975, 1999.

18. W. Chen and C. Peng, "Purification of Protease from *Aeromonas* and Using as Immunization Factor," *Appl. Microbiol. Biotechnol.*, vol. 25, pp. 30-35, 2003.
19. S. Sh. Omer and H. G. Humadi, "Qualitative and Quantitative Screening of Alkaline Protease Production from Some Pathogenic Bacteria," *Journal of Kerbala University*, vol. 11, no. 3, pp. 309-314, 2013.
20. C. Cupp-Enyard, "Sigma's a Specific Protease Activity Assay Casein as a Substrate," *Journal of Visual Experiments*, vol. 19, pp. 1-2, 2013.
21. M. C. Wheeler, M. H. Roe, F. I. Kaplan, P. M. Schlivert, and J. K. Todd, "Outbreak of Group A *Streptococcus* Septicemia in Children: Clinical, Epidemiologic, and Microbiological Correlates," *JAMA*, vol. 266, pp. 533-537, 1991.