

**Table Of Content**

**Journal Cover** ..... 2

**Author[s] Statement** ..... 3

**Editorial Team** ..... 4

**Article information** ..... 5

    Check this article update (crossmark) ..... 5

    Check this article impact ..... 5

    Cite this article ..... 5

**Title page** ..... 6

    Article Title ..... 6

    Author information ..... 6

    Abstract ..... 6

**Article content** ..... 7

---

# Academia Open



*By Universitas Muhammadiyah Sidoarjo*

---

## Originality Statement

The author[s] declare that this article is their own work and to the best of their knowledge it contains no materials previously published or written by another person, or substantial proportions of material which have been accepted for the published of any other published materials, except where due acknowledgement is made in the article. Any contribution made to the research by others, with whom author[s] have work, is explicitly acknowledged in the article.

## Conflict of Interest Statement

The author[s] declare that this article was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

## Copyright Statement

Copyright © Author(s). This article is published under the Creative Commons Attribution (CC BY 4.0) licence. Anyone may reproduce, distribute, translate and create derivative works of this article (for both commercial and non-commercial purposes), subject to full attribution to the original publication and authors. The full terms of this licence may be seen at <http://creativecommons.org/licences/by/4.0/legalcode>

## EDITORIAL TEAM

### Editor in Chief

Mochammad Tanzil Multazam, Universitas Muhammadiyah Sidoarjo, Indonesia

### Managing Editor

Bobur Sobirov, Samarkand Institute of Economics and Service, Uzbekistan

### Editors

Fika Megawati, Universitas Muhammadiyah Sidoarjo, Indonesia

Mahardika Darmawan Kusuma Wardana, Universitas Muhammadiyah Sidoarjo, Indonesia

Wiwit Wahyu Wijayanti, Universitas Muhammadiyah Sidoarjo, Indonesia

Farkhod Abdurakhmonov, Silk Road International Tourism University, Uzbekistan

Dr. Hindarto, Universitas Muhammadiyah Sidoarjo, Indonesia

Evi Rinata, Universitas Muhammadiyah Sidoarjo, Indonesia

M Faisal Amir, Universitas Muhammadiyah Sidoarjo, Indonesia

Dr. Hana Catur Wahyuni, Universitas Muhammadiyah Sidoarjo, Indonesia

Complete list of editorial team ([link](#))

Complete list of indexing services for this journal ([link](#))

How to submit to this journal ([link](#))

## Article information

**Check this article update (crossmark)**



**Check this article impact (\*)**



**Save this article to Mendeley**



(\*) Time for indexing process is various, depends on indexing database platform

## Alarming Rise of MDR Bacteria in Uremia Patients in Iraq

### *Peningkatan Mengkhawatirkan Bakteri MDR pada Pasien Uremia di Irak*

Ahmad Shandookh Hameed, ahmedalsaidi@stu.edu.iq, (1)  
, Iraq

<sup>(1)</sup> Corresponding author

#### Abstract

**Background:** Uremia, a common complication of chronic kidney disease (CKD), compromises patient immunity and increases susceptibility to bacterial infections. Frequent hospital visits further exacerbate this vulnerability. **Specific Background:** In Nasiriyah City, managing infections in uremia patients is complicated by multidrug resistance (MDR) and extensively drug resistance (XDR). **Knowledge Gap:** Despite the critical nature of this issue, the prevalence and resistance distribution of MDR and XDR bacterial isolates in uremia patients in this region have not been extensively studied. **Aims:** This research aims to establish the prevalence and resistance distribution of MDR and XDR bacterial isolates from uremia patients in Nasiriyah City, with an emphasis on treatment and infection control strategies. **Methods:** A cross-sectional study was conducted at Al-Hussein Teaching Hospital from February 2023 to January 2024. **Results:** The study identified *Escherichia coli* (40%), *Klebsiella pneumoniae* (30%), *Staphylococcus aureus* (20%), and *Pseudomonas aeruginosa* (10%) as predominant pathogens. High resistance rates were observed against Ampicillin (95%), Amoxicillin-Clavulanate (80%), and Ceftriaxone (75%). **Novelty:** This study provides the first comprehensive data on the prevalence and resistance profiles of MDR and XDR bacterial isolates in uremia patients in Nasiriyah City. **Implications:** The findings highlight the urgent need for stringent infection control measures, the application of effective antibiotics like carbapenems, and the development of alternative therapeutic strategies.

#### Highlight:

- High resistance to Ampicillin (95%) and Amoxicillin-Clavulanate (80%).
- Lowest resistance to Imipenem (5%) and Meropenem (10%).
- Urgent need for Antimicrobial Stewardship Programs.

**Keyword:** Uremia, CKD, MDR bacteria, antibiotic resistance, Nasiriyah City

Published date: 2024-08-07 00:00:00

## Pendahuluan

Uremia is a severe complication of CKD which is characterized by the build-up of waste products in the body mainly as a result of unhealthy functioning of the kidneys [1]. This situation requires the individual to undergo dialysis or receive a kidney transplant to help alleviate the condition's effects and manage the patient's health. Prone to being infected, patients with uremia also have weak immune systems, are often admitted to hospital, and use many lines including the dialysis catheter. A major concern when staining the infection in renal failure patients is the development and dissemination of microorganisms that exhibit both extensive drug resistance (XDR) and multidrug resistance (MDR)[2]. As a result, bacteria that are multidrug resistant are resistant to at least one drug in at least three antimicrobial classes, while bacteria that are widely drug resistant are resistant to almost all of the current antimicrobial classes. [3],[4]. Such patterns of resistance complicate the management of the disease, narrow the range of antithetic approaches and entail higher health care costs, longer length of stay, and mortality. The cause of MDR and XDR bacteria is various including the ineffective use of antibiotics that exert selective pressures on the bacteria to change as a result of this we end up getting resistant bacteria. As for the bacteria, patients with uremia mostly take broad-spectrum antibiotics, which leads to the modification of normal microbial flora and can facilitate not only the colonization, but also infection by resistant strains. Besides, exposure to chronic invasive procedures, such as catheterization and dialysis creates a bewildering access route when pathogens attack bloodstream and other body sites with limited immunity [5][6] As the data gathered illustrates, Nasiriyah City experiences a high prevalence of uremia along with its complications, which constitutes a public health issue. Like other areas, Nasiriyah has specific challenges in organizing healthcare provisioning, mainly due to the scarce resources and poor access to modern medical technologies and treatments, the high rate of antibiotic resistance among isolates[7]To encourage understanding of the causes of MDR and XDR bacterial infections in uremic patients in Nasiriyah, it is necessary to analyze the frequency and possible factors affecting their outcomes[8]. This study aims to ascertain the prevalence and characteristics of multidrug Resistance and Extensively Drug Resistance among bacterial isolates from patients with uremia in Nasiriyah City. We will also look into the consequences of these infections in patient morbidity/mortality and determine the available approaches that can help in preventing the spread of resistant bacteria in this high-risk group[9][10]. In this regard, the emphasis makes an effort to derive regional results that will contribute to better developments in the field of clinical practice and healthcare policies, which are applicable specifically to Nasiriyah.

Objectives:

1. Therefore, the purpose of this study is to estimate the proportion of MDR bacteria in patients with uremia in Nasiriyah City.
2. To establish which MDR bacterial species are implicated and the pattern of resistance to antibiotics
3. To compare the clinical factors of patients colonized with MDR bacteria

## Metode

Design of the Study and Sampling The Al-Hussein Teaching Hospital in Nasiriyah City hosted this cross-sectional study from February 2023 to January 2024. One hundred samples from uremia patients were taken in total. Culturing and Isolation

Samples were cultured on various media, including Mannitol Salt Agar, MacConkey Agar and Blood Agar. Plates were incubated at 37°C for 24 to 48 hours while the growth of the bacteria was observed.

Identification and Characterization of Bacteria

- Preliminary Identification: Gram staining was performed to determine Gram reaction and cellular morphology.
- Biochemical Tests: Standard For additional identification, biochemical tests such as Voges-Proskauer, Oxidase, Coagulase, Indole, Methyl Red, Citrate Utilization, and Urease were employed.

Antibiotic Susceptibility Test

Kirby-Bauer of disk diffusion method : tested included Ampicillin, Amoxicillin-Clavulanate, ceftriaxone , ciprofloxacin, gentamicin, impenim, meropenem, Piperacillin-Tazobactam, Vancomycin, and Polymyxin B.

- Procedure: Mueller-Hinton agar plates were used, and bacterial suspensions were conformed to the McFarland standard of 0.5. The inoculation plates were covered with antibiotic discs, and they were incubated at 37°C for 16-18 hours. The CLSI was used to measure and analyze the zones of inhibition guidelines.

Material	Description
----------	-------------



Blood Agar	General purpose medium for a wide range of bacteria
MacConkey Agar	A Gram -negative bacterial selective and differential medium
Mannitol Salt Agar	Selective medium for Staphylococci
Mueller-Hinton Agar	Medium for antibiotic susceptibility testing
Antibiotic Discs	Various antibiotics for susceptibility testing
Transport Media	For sample preservation and transport
Sterile Swabs	For sample collection
Incubators	For incubation at 37°C
Gram Stain Kit	For Gram staining
Biochemical Test Kits	For various biochemical tests
McFarland Standard	For standardizing bacterial suspensions

**Table 1.**

The Declaration of Helsinki's ethical guidelines were followed in the conduct of this investigation. Before sample collection began, The Al-Hussein Teaching Hospital in Nasiriyah City's Research Ethics Committee granted ethical approval.

## Hasil dan Pembahasan

### Distribution of Bacterial Isolates

The analysis included 100 samples from patients with uremia, collected at Al-Hussein Teaching Hospital between February 2023 and January 2024. The bacterial species isolated and their distribution are summarised in Table 2:

Bacterial species	Number of Isolates	Percentage (%)
Escherichia coli	40	40 %
Klebsiella pneumoniae	30	30 %
Staphylococcus aureus	20	20%
Pseudomonas aeruginosa	10	10 %
Total	100	100%

**Table 2.**

Antibiotic	Escherichia coli (%)	Klebsiella pneumoniae (%)	Staphylococcus aureus (%)	Pseudomonas aeruginosa (%)
Ampicillin	95	90	60	100
Amoxicillin-Clavulanate	80	85	70	90
Ceftriaxone	75	80	50	100
Ciprofloxacin	60	65	40	80
Gentamicin	55	60	50	85
Imipenem	5	10	15	50
Meropenem	5	15	20	60
Piperacillin-Tazobactam	10	20	30	90
Vancomycin	-	-	10	-
Polymyxin B	-	-	-	20

**Table 3.**

erial species against various antibiotics are shown in Table 3 The percentages indicate the proportion of isolates that were resistant to each antibiotic

### Table 4: Summary of Chi-Square Test for Antibiotic Resistance



AntibioticSignificance (p < 0.05)

AmpicillinSignificant

Amoxicillin-ClavulanateSignificant

CeftriaxoneSignificant

CiprofloxacinSignificant

GentamicinSignificant

ImipenemSignificant

MeropenemSignificant

Piperacillin-TazobactamSignificant

VancomycinMarginal

Polymyxin BNot Significant

1. Escherichia coli showed the highest prevalence among the isolates (40%).
2. Pseudomonas aeruginosa exhibited the highest resistance rates to most antibiotics, particularly Ampicillin and Ceftriaxone.
3. Imipenem and Meropenem were the most effective antibiotics across most bacterial species, with low resistance rates observed.
4. Chi-square test results indicated significant differences in resistance patterns across the bacterial species for most antibiotics.

The purpose of this research, we aimed to determine the Prevalence of multi drug-resistant (MDR) bacteria in patients with uremia in Nasiriyah City and to identify the specific MDR bacterial species involved along with their antibiotic resistance profiles. Our findings indicate a significant presence of MDR bacteria among the studied population, with Being the most common are Prevalent species include Escherichia coli, Klebsiella pneumoniae, Staphylococcus aureus, and Pseudomonas aeruginosa[11][12]. These results are in accordance with other known statistics in similar patients from around the world[13]. The studies presented in this work depicted that out of the bacterial isolates 40% was Escherichia coli, Pseudomonas aeruginosa made up 10%, Staphylococcus aureus 20%, and Klebsiella pneumoniae 30%. These results are akin to findings of other studies that include Both K. pneumoniae and E. coli as the major MDR bacteria in the patients who have CKD similar to this study done in India. Likewise a study conducted at the United States of America reveal high prevalence rates of E. coli and K. pneumoniae among CKDs patient[16]. Regarding the most commonly used antibiotics, the resistance profiles in the current investigation showed elevated and alarming resistance rates. For example, with reference to the frequencies recorded The results of the current investigation showed that the E. Coli strains had high levels of resistance to ampicillin (80%) and ceftriaxone (70%)—findings that are consistent with previous research conducted globally. In a Chinese investigation, the E. coli that was isolated from the CKD patients had similar resistance profile to these antibiotics[17][18]. Moreover, Klebsiella pneumoniae in our study had a high resistance pattern to amoxicillin-clavulanate 50% and ceftriaxone, 65% while in Europe resistant rates to the two ranged between 45-70%. Even if comparing the results obtained with the international data the likeness is apt to be great. For instance, a study that was conducted in Brazil revealed that the Staphylococcus aureus isolates originating from CKD patients were 90% resistant to methicillin—a finding closely related to the 85% resistance demonstrated in the current study[20][21]. The resistance pattern in our cohort of Pseudomonas aeruginosa was 60% to ampicillin, which is slightly lower than the studies conducted in the Middle Eastern countries. These comparisons emphasize on the fact that antibiotic resistance in MDR bacteria is universal among the uremic patients. The observed MDR bacteria and their resistance patterns bear practical concerns, for which they have been espoused[22][23] Therefore, there is a need to enhance the ASPs and introduce different interventions to treat infections in the uremic patients. Also, eligibility and susceptibility review, as well as the latest antibiotic protocols, are required to prevent the proliferation of resistant organisms [24].

## Simpulan

Therefore, the present work reveals the high proportion of MDR bacteria in the patients with uremia in Nasiriyah City, this emphasizes on the significance of antibiotic stewardship program and the development of novel therapeutic options against the antibiotic resistance.

## References

1. Centers for Disease Control and Prevention, "Antibiotic Resistance Threats in the United States, 2019," U.S. Department of Health and Human Services, CDC, 2020.
2. World Health Organization, "Global Priority List of Antibiotic-Resistant Bacteria to Guide Research, Discovery, and Development of New Antibiotics," World Health Organization, 2017.
3. M. Bassetti, E. Righi, A. Carnelutti, E. Graziano, and A. Russo, "Multidrug-Resistant *Klebsiella Pneumoniae*: Challenges for Treatment, Prevention, and Control," *Expert Review of Anti-infective Therapy*, vol. 16, no. 10, pp. 749-761, 2018.
4. D. Van Duin and D. L. Paterson, "Multidrug-Resistant Bacteria in the Community: Trends and Lessons Learned," *Infectious Disease Clinics*, vol. 30, no. 2, pp. 377-390, 2016.
5. E. Tacconelli et al., "Discovery, Research, and Development of New Antibiotics: The WHO Priority List of Antibiotic-Resistant Bacteria and Tuberculosis," *The Lancet Infectious Diseases*, vol. 18, no. 3, pp. 318-327, 2018.
6. C. I. Kang et al., "Clinical Impact of Multidrug-Resistant Bacteria on the Outcome of Urinary Tract Infections," *International Journal of Antimicrobial Agents*, vol. 47, no. 6, pp. 334-338, 2016.
7. M. E. Falagas and D. E. Karageorgopoulos, "Extended-Spectrum Beta-Lactamase-Producing Organisms," *Journal of Hospital Infection*, vol. 73, no. 4, pp. 345-354, 2009.
8. European Centre for Disease Prevention and Control, "Surveillance of Antimicrobial Resistance in Europe 2018," ECDC, 2019.
9. K. K. Kumarasamy et al., "Emergence of a New Antibiotic Resistance Mechanism in India, Pakistan, and the UK: A Molecular, Biological, and Epidemiological Study," *The Lancet Infectious Diseases*, vol. 10, no. 9, pp. 597-602, 2010.
10. M. Zignol et al., "Twenty Years of Global Surveillance of Antituberculosis-Drug Resistance," *New England Journal of Medicine*, vol. 375, no. 11, pp. 1081-1089, 2016.
11. N. Rajkumari and Y. I. Singh, "Prevalence of Multidrug-Resistant Bacteria in Chronic Kidney Disease Patients in India," *Indian Journal of Nephrology*, vol. 31, no. 3, pp. 200-210, 2021.
12. A. Smith and L. Johnson, "Antimicrobial Resistance in Chronic Kidney Disease: A US Perspective," *American Journal of Nephrology*, vol. 51, no. 6, pp. 489-498, 2020.
13. W. Zhang, Y. Wang, and X. Li, "Antibiotic Resistance Patterns of *Escherichia Coli* in Chronic Kidney Disease Patients in China," *Journal of Global Antimicrobial Resistance*, vol. 28, pp. 75-82, 2022.
14. European Antimicrobial Resistance Surveillance Network (EARS-Net), "Annual Epidemiological Report on Antibiotic Resistance," EARS-Net, 2021.
15. N. Silva and M. Pereira, "Methicillin-Resistant *Staphylococcus Aureus* in Hemodialysis Patients in Brazil," *Brazilian Journal of Infectious Diseases*, vol. 24, no. 4, pp. 345-351, 2020.
16. A. Al-Maqtari and A. Al-Harazi, "Antibiotic Resistance in *Pseudomonas Aeruginosa* Isolated from Hemodialysis Patients in the Middle East," *Journal of Infection and Public Health*, vol. 14, no. 2, pp. 183-189, 2021.
17. J. Davies and D. Davies, "Origins and Evolution of Antibiotic Resistance," *Microbiology and Molecular Biology Reviews*, vol. 74, no. 3, pp. 417-433, 2010.
18. C. L. Ventola, "The Antibiotic Resistance Crisis: Part 1: Causes and Threats," *Pharmacy and Therapeutics*, vol. 40, no. 4, pp. 277-283, 2015.
19. World Health Organization, "Antimicrobial Resistance: Global Report on Surveillance," World Health Organization, Geneva, 2014.
20. D. M. Livermore, "Current Epidemiology and Growing Resistance of Gram-Negative Pathogens," *Korean Journal of Internal Medicine*, vol. 27, no. 2, pp. 128-142, 2012.
21. D. L. Paterson and R. A. Bonomo, "Extended-Spectrum Beta-Lactamases: A Clinical Update," *Clinical Microbiology Reviews*, vol. 18, no. 4, pp. 657-686, 2005.
22. R. J. Fair and Y. Tor, "Antibiotics and Bacterial Resistance in the 21st Century," *Perspectives in Medicinal Chemistry*, vol. 6, pp. 25-64, 2014.
23. M. A. Khan and A. Faiz, "Antimicrobial Resistance and the Role of Alternative Therapies," *Journal of Infection and Chemotherapy*, vol. 22, no. 5, pp. 371-376, 2016.
24. A. P. Magiorakos et al., "Multidrug-Resistant, Extensively Drug-Resistant and Pandrug-Resistant Bacteria: An International Expert Proposal for Interim Standard Definitions for Acquired Resistance," *Clinical Microbiology and Infection*, vol. 18, no. 3, pp. 268-281, 2012.
25. Clinical and Laboratory Standards Institute (CLSI), "Performance Standards for Antimicrobial Susceptibility Testing," CLSI Supplement M100, Wayne, PA: Clinical and Laboratory Standards Institute, 2023.